

Mathematics in Art and Architecture GEK1518K


Helmer Aslaksen
Department of Mathematics
National University of Singapore
aslaksen@math.nus.edu.sg
www.math.nus.edu.sg/aslaksen/

The Golden Ratio

The Golden Ratio

Suppose we want to divide a line AB at a point C such that AC is longer than BC and the ratio of AC to BC is equal to the ratio of AB to AC. In other words,

$$\text{whole} / \text{long} = \text{long} / \text{short}.$$



We will determine the value of this common ratio $\varphi = \frac{AC}{BC} = \frac{AB}{AC}$.

If $CB = a$, then $AC = a\varphi$ and $AB = a\varphi^2$. Hence,


$$\begin{aligned} a\varphi^2 &= a\varphi + a \\ \varphi^2 &= \varphi + 1 \end{aligned}$$

The positive solution is $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$. This value of φ is known as the *golden ratio*. It is also known as the *golden section* or the *divine ratio*.

Conversely, given a line segment AB, we may want to extend it to a line AE such that

$\frac{AE}{AB} = \varphi$. We will call this a golden ration extension. To construct this extension

geometrically, start with a line AB of length a and construct a square ABCD. Bisect AB and let the midpoint be M. With M as a center, draw an arc with radius MC to cut AB produced at E. Let us now check the ratios AE/AB and AB/BE. To do so, we need to find the lengths MC and AE.



By Pythagoras Theorem,

$$\begin{aligned} (MC)^2 &= (MB)^2 + (BC)^2 \\ &= \left(\frac{a}{2}\right)^2 + a^2 \\ &= \left(\frac{5}{4}\right) a^2 \\ MC &= \left(\frac{\sqrt{5}}{2}\right) a \end{aligned}$$

$$\begin{aligned} AE &= AM + ME \\ &= AM + MC \\ &= \left(\frac{1}{2}\right) a + \left(\frac{\sqrt{5}}{2}\right) a \\ &= a (1 + \sqrt{5})/2 \end{aligned}$$

Hence

$$\frac{AE}{AB} = \frac{a(1 + \sqrt{5}/2)}{a} = (1 + \sqrt{5})/2,$$


and since

$$\frac{AB}{BE} = \frac{a}{a(\sqrt{5}/2 - 1/2)} = (1 + \sqrt{5})/2,$$

we see that $\frac{AE}{AB} = \frac{AB}{BE} = \phi$.

Observe that the rectangle AEFD has length $a \phi$ and height a . We call a rectangle with dimensions a by $a \phi$ a *golden rectangle*.


If the short segment has length 1, then the long segment has length ϕ and the whole segment has length ϕ^2 . If the long segment has length 1, then the short segment has length $1/\phi$ and the whole segment has length ϕ . If the whole segment has length 1, then the long segment has length $1/\phi$ and the short segment has length $1/\phi^2$.



Extending line AB to ABE by a length of a/ϕ to get a golden ratio extension

The above construction shows how to get a golden ratio extension of AB to ABE. We now want to construct a golden ratio cut, that is, start with AB and find a point C such that $AB/AC=AC/CB = \phi$. It turns out that we have done all the hard work already. In the above golden ratio extension, the length of BE is a/ϕ . But this is exactly the length of the long piece in a golden ratio cut. Using a compass centered at B with radius $BE = a/\phi$, cut the line AB at X. So, $BX = BE = a/\phi$. Notice that AX is the smaller part while BX is the larger part of the line segment because $BX = a/\phi > \frac{1}{2} a$. Since $\frac{AB}{BE} = \phi$ and $BX = BE$,

$$\frac{AB}{BX} = \frac{AB}{BE} = \phi.$$




Cutting a given line AB into 2 parts to achieve the golden ratio

Now we know how to do both a golden ratio extension and a golden ratio cut.

Constructing a Regular Pentagon using the Golden Ratio

We will use the golden ratio to help us construct a regular pentagon. Observe that if we divide a regular pentagon ABCDE into 5 isosceles triangles as shown in Figure 5.2.1, then, for each isosceles triangle, the interior angle at F is 72° .




A regular pentagon

We will construct a regular pentagon within a circumscribing circle. To do that, we need a way to construct the angle of 72° . Euclid came up with the following method to get this angle.

- 1) Begin with a line ABE, which is divided into the golden ratio at B, with AB longer than BE.
- 2) Using compasses centered at B and E and of radius AB, draw two arcs of circles to intersect at point C. So $AB = BC = CE$ and we will prove later that $AC = AE$.
- 3) Form the triangle CEB. Since $BC = CE$, so $\angle CBE = \angle CEB$. Let these two angles be α . So $\angle BCE = 180^\circ - 2\alpha$.
- 4) Consider the triangle ACE. Since $AC = AE$, $\angle ACE = \angle CEB = \alpha$. So $\angle CAB = 180^\circ - 2\alpha$.

- 5) Consider the triangle ACB. $\angle ACB = \angle ACE - \angle BCE = 3\alpha - 180^\circ$.
 Since $AB = BC$, $\angle CAB = 3\alpha - 180^\circ$.
- 6) From (4) and (5), $\angle CAB = 3\alpha - 180^\circ = 180^\circ - 2\alpha$. Solving, we get $\alpha = 72^\circ$.

Now that we obtain the angle $\alpha = 72^\circ$ by the above construction, construct a circle at center E, radius less than BE. The circle cuts BE and CE at X and Y respectively. X and Y form two vertices of our regular pentagon. We can get the remaining three vertices of the pentagon using compasses with radius XY and cutting the circle.





Now, we prove that $AC = AE$. Let the length of $AB = a$ and so, $BE = a/\phi$. So $AE = a(1 + 1/\phi)$. By Equation (5.1), $AE = a\phi$. So we want to prove that $AC = a\phi$.


Find the mid-point N of BE and the line CN is perpendicular to BE . We apply Pythagoras Theorem to get AC .




$$\begin{aligned}
 (AC)^2 &= CN^2 + AN^2 \\
 &= (CE^2 - NE^2) + (AB + BN)^2 \\
 &= (a^2 - (a/2\phi)^2) + (a + (a/2\phi))^2 \\
 &= a^2(2 + 1/\phi) \\
 &= a^2(1 + \phi) \quad \text{by Equation (5.1)} \\
 &= a^2\phi^2 \quad \text{by Equation (5.2)}
 \end{aligned}$$

Therefore, $AC = a\phi = AE$.




THE GOLDEN SECTION





Battlement
A, Merlon; B, Crenel;
C, Loophole; D, Machicolation



Comparison of (left) ground plan of a medieval castle with circular towers, and (right) a bastioned plan, showing how the new design eliminated the dead ground (shaded) at the foot of the tower.

