

AN ALGORITHM FOR CONSTRUCTING MULTIDIMENSIONAL BIORTHOGONAL PERIODIC MULTIWAVELETS

SAY SONG GOH AND K. M. TEO

ABSTRACT. This paper deals with the problem of constructing multidimensional biorthogonal periodic multiwavelets from a given pair of biorthogonal periodic multiresolutions. Biorthogonal polyphase splines are introduced to reduce the problem to a matrix extension problem, and an algorithm for solving the matrix extension problem is derived. Sufficient conditions for collections of periodic multiwavelets to form a pair of biorthogonal Riesz bases of the entire function space are also obtained.

1. INTRODUCTION

One of the important problems in wavelet analysis is the construction of biorthogonal wavelets or multiwavelets from a given pair of biorthogonal multiresolutions. Here, we consider this problem for the space $L^2([0, 2\pi)^s)$, $s \geq 1$, of 2π -periodic square-integrable complex-valued functions over $[0, 2\pi)^s$.

For the space $L^2(\mathbb{R})$, it is well known that biorthogonal wavelets with dilation 2 can be constructed from multiresolutions generated by biorthogonal scaling functions with the use of simple alternating flip formulas. The problem of constructing biorthogonal multiwavelets is much more complicated. The paper [9] contains an algorithmic approach for the construction of biorthogonal multiwavelets with arbitrary dilation from multiresolutions generated by several biorthogonal scaling functions. However, such an algorithm is unavailable for $L^2(\mathbb{R}^s)$, $s > 1$, with arbitrary dilation matrices.

This paper presents such an algorithm for the space $L^2([0, 2\pi)^s)$, $s \geq 1$. Periodic wavelets and multiwavelets were studied in one or higher dimensions in [1], [2], [4], [8], [11], [12], [13], but there is still no work done in the biorthogonal setting. To fix

1991 *Mathematics Subject Classification.* 41A63, 41A65, 42C15.

notations, let M be an $s \times s$ matrix with integer entries such that all its eigenvalues lie outside the unit circle. For $k \geq 0$, let \mathcal{L}_k denote a full collection of coset representatives of $\mathbb{Z}^s / M^k \mathbb{Z}^s$. Then $\mathbb{Z}^s = \bigcup_{\ell \in \mathcal{L}_k} (\ell + M^k \mathbb{Z}^s)$, and for any distinct $\ell_1, \ell_2 \in \mathcal{L}_k$, $(\ell_1 + M^k \mathbb{Z}^s) \cap (\ell_2 + M^k \mathbb{Z}^s) = \emptyset$. Similarly, we let \mathcal{R}_k denote a full collection of coset representatives of $\mathbb{Z}^s / D^k \mathbb{Z}^s$, where $D = M^T$.

For $k \geq 0$, consider $\mathcal{S}(M^k)^{r \times \rho}$, the class of periodic sequences of $r \times \rho$ complex matrices of period M^k , that is, $H_k(\ell + M^k p) = H_k(\ell)$ for all $H_k \in \mathcal{S}(M^k)^{r \times \rho}$, $\ell, p \in \mathbb{Z}^s$. For $H_k \in \mathcal{S}(M^k)^{r \times \rho}$, we define its finite Fourier transform by

$$\widehat{H}_k(j) = \sum_{\ell \in \mathcal{L}_k} H_k(\ell) e^{-ij \cdot (2\pi M^{-k} \ell)}, \quad j \in \mathcal{R}_k.$$

Then $\widehat{H}_k \in \mathcal{S}(D^k)^{r \times \rho}$, and the sequence H_k can be recovered from \widehat{H}_k by

$$H_k(\ell) = \frac{1}{|\mathcal{R}_k|} \sum_{j \in \mathcal{R}_k} \widehat{H}_k(j) e^{i\ell \cdot (2\pi D^{-k} j)}, \quad \ell \in \mathcal{L}_k.$$

In addition,

$$\frac{1}{|\mathcal{R}_k|} \sum_{j \in \mathcal{R}_k} |\widehat{H}_k(j)|^2 = \sum_{\ell \in \mathcal{L}_k} |H_k(\ell)|^2, \quad H_k \in \mathcal{S}(M^k), \quad (1.1)$$

where $\mathcal{S}(M^k) := \mathcal{S}(M^k)^{1 \times 1}$.

We say that a sequence of subspaces $\{V_k : k \geq 0\}$ of $L^2([0, 2\pi]^s)$ is a *periodic multiresolution of $L^2([0, 2\pi]^s)$ with multiplicity r and dilation matrix M* if it satisfies the following conditions:

- MR1 For $k = 0, 1, \dots$, $\dim V_k = r |\det(M^k)|$ and there exist functions $\phi_k^m \in V_k$, $m = 1, 2, \dots, r$, such that $\{T_k^\ell \phi_k^m : m = 1, 2, \dots, r, \ell \in \mathcal{L}_k\}$ is a basis for V_k , where $T_k^\ell f := f(\cdot - 2\pi M^{-k} \ell)$.
- MR2 For $k = 0, 1, \dots$, $V_k \subseteq V_{k+1}$.
- MR3 $\overline{\bigcup_{k \geq 0} V_k} = L^2([0, 2\pi]^s)$.

The functions ϕ_k^m , $k \geq 0$, $m = 1, 2, \dots, r$, are called *scaling functions*. Suppose that $\{\widetilde{V}_k : k \geq 0\}$ is another periodic multiresolution of $L^2([0, 2\pi]^s)$ with multiplicity r and dilation matrix M generated by scaling functions $\widetilde{\phi}_k^m$, $k \geq 0$, $m = 1, 2, \dots, r$. The two multiresolutions $\{V_k : k \geq 0\}$ and $\{\widetilde{V}_k : k \geq 0\}$ are said to be *biorthogonal* if

the collections $\{T_k^\ell \phi_k^m : m = 1, 2, \dots, r, \ell \in \mathcal{L}_k\}$ and $\{T_k^\ell \tilde{\phi}_k^m : m = 1, 2, \dots, r, \ell \in \mathcal{L}_k\}$ are biorthogonal for every $k \geq 0$, that is,

$$\langle T_k^j \phi_k^m, T_k^\ell \tilde{\phi}_k^\mu \rangle = \delta_{j\ell} \delta_{m\mu}, \quad k \geq 0, \quad j, \ell \in \mathcal{L}_k, \quad m, \mu = 1, 2, \dots, r. \quad (1.2)$$

The scaling functions ϕ_k^m and $\tilde{\phi}_k^m$, $k \geq 0$, $m = 1, 2, \dots, r$, satisfy the *periodic matrix refinement equations*

$$\phi_k = \sum_{\ell \in \mathcal{L}_{k+1}} H_{k+1}(\ell) T_{k+1}^\ell \phi_{k+1}, \quad \tilde{\phi}_k = \sum_{\ell \in \mathcal{L}_{k+1}} \tilde{H}_{k+1}(\ell) T_{k+1}^\ell \tilde{\phi}_{k+1}, \quad (1.3)$$

where $\phi_k := (\phi_k^1, \dots, \phi_k^r)^T$, $\tilde{\phi}_k := (\tilde{\phi}_k^1, \dots, \tilde{\phi}_k^r)^T$, and $H_{k+1}, \tilde{H}_{k+1} \in \mathcal{S}(M^{k+1})^{r \times r}$, for every $k \geq 0$. Writing in terms of Fourier coefficients, (1.3) is equivalent to

$$\hat{\phi}_k(n) = \hat{H}_{k+1}(n) \hat{\phi}_{k+1}(n), \quad \hat{\tilde{\phi}}_k(n) = \hat{\tilde{H}}_{k+1}(n) \hat{\tilde{\phi}}_{k+1}(n),$$

for all $n \in \mathbb{Z}^s$.

In this paper, we seek functions $\psi_k^m, \tilde{\psi}_k^m \in L^2([0, 2\pi]^s)$, $k \geq 0$, $m = 1, 2, \dots, r(d-1)$, where $d = |\det(M)|$, called *multiwavelets* satisfying the biorthogonality relation

$$\langle T_k^j \psi_k^m, T_k^\ell \tilde{\psi}_k^\mu \rangle = \delta_{j\ell} \delta_{m\mu}, \quad (1.4)$$

for $k \geq 0$, $m, \mu = 1, 2, \dots, r(d-1)$, $j, \ell \in \mathcal{L}_k$, with the sets

$$W_k := \langle \{T_k^\ell \psi_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\} \rangle \quad (1.5)$$

and

$$\tilde{W}_k := \langle \{T_k^\ell \tilde{\psi}_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\} \rangle \quad (1.6)$$

satisfying the direct sums

$$V_{k+1} = V_k \oplus W_k, \quad \tilde{V}_{k+1} = \tilde{V}_k \oplus \tilde{W}_k, \quad (1.7)$$

and the orthogonality properties

$$W_k \perp \tilde{V}_k, \quad \tilde{W}_k \perp V_k, \quad (1.8)$$

for $k \geq 0$. Since $\dim V_k = rd^k$, (1.7) will imply that each of the collections $\{T_k^\ell \psi_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\}$ and $\{T_k^\ell \tilde{\psi}_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\}$ is linearly independent.

In [8], a polyphase spline approach was proposed for the study of multiresolutions and multiwavelets in $L^2([0, 2\pi]^s)$ for the orthonormal and semi-orthogonal settings. Here, we extend it to the biorthogonal setting. In Section 2, we characterize the biorthogonality relation (1.2) in terms of polyphase splines. In Section 3, this characterization is used to reduce the biorthogonal multiwavelet construction problem to a matrix extension problem, and a constructive solution is provided. The paper concludes in Section 4 with sufficient conditions for collections of biorthogonal multiwavelets to form a pair of biorthogonal Riesz bases of $L^2([0, 2\pi]^s)$.

2. CHARACTERIZATION OF BIORTHOGONALITY

For $k \geq 0$, let $\phi_k^m, \tilde{\phi}_k^m$, $m = 1, 2, \dots, r$, be functions in $L^2([0, 2\pi]^s)$. Consider the sets $S_k := \{T_k^\ell \phi_k^m : m = 1, 2, \dots, r, \ell \in \mathcal{L}_k\}$ and $\tilde{S}_k := \{T_k^\ell \tilde{\phi}_k^m : m = 1, 2, \dots, r, \ell \in \mathcal{L}_k\}$. Define *polyphase splines* $v_{k,j}^m$ in $L^2([0, 2\pi]^s)$ by

$$v_{k,j}^m(x) := \sum_{p \in \mathbb{Z}^s} \hat{\phi}_k^m(j + D^k p) e^{i(j + D^k p) \cdot x},$$

for $m = 1, 2, \dots, r$, $j \in \mathcal{R}_k$, where $\hat{\phi}_k^m(n)$ denote the Fourier coefficients of ϕ_k^m . Similarly, we define $\tilde{v}_{k,j}^m$ using $\hat{\tilde{\phi}}_k^m(n)$, the Fourier coefficients of $\tilde{\phi}_k^m$.

The polyphase splines $v_{k,j}^m$ were first introduced in [8], and they were used to characterize the linear independence of the set S_k . Here, we shall use the two collections of polyphase splines $\{v_{k,j}^m : j \in \mathcal{R}_k, m = 1, 2, \dots, r\}$ and $\{\tilde{v}_{k,j}^m : j \in \mathcal{R}_k, m = 1, 2, \dots, r\}$ to characterize the biorthogonality of S_k and \tilde{S}_k .

Theorem 2.1. *For $k \geq 0$, the sets S_k and \tilde{S}_k are biorthogonal, that is,*

$$\langle T_k^p \phi_k^m, T_k^q \tilde{\phi}_k^\mu \rangle = \delta_{pq} \delta_{m\mu}, \quad p, q \in \mathcal{L}_k, \quad m, \mu = 1, 2, \dots, r,$$

if and only if

$$\left(\langle v_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle \right)_{m,\mu=1}^r = \frac{1}{d^k} I_r \quad j \in \mathcal{R}_k,$$

where I_r denotes the $r \times r$ identity matrix.

Proof. First, we note that it suffices to establish the theorem for any chosen collections of coset representatives \mathcal{L}_k and \mathcal{R}_k . Thus we shall select the particular

collections of \mathcal{L}_k and \mathcal{R}_k described in [8, Section 3]. The procedure is as follows. Choose bases $\{e'_1, \dots, e'_s\}$ and $\{e''_1, \dots, e''_s\}$ of the free abelian group \mathbb{Z}^s , and sets of generators $\{f'_1, \dots, f'_s\}$ and $\{g'_1, \dots, g'_s\}$ of $M^k\mathbb{Z}^s$ and $D^k\mathbb{Z}^s$ respectively, such that

$$f'_i = n_i e'_i, \quad g'_i = n_i e''_i, \quad i = 1, 2, \dots, s.$$

Here, n_1, \dots, n_s are positive integers such that

$$d^k = |\det(M^k)| = |\det(D^k)| = n_1 n_2 \cdots n_s.$$

Thus, if we set

$$\mathcal{L}_k = \{m_1 e'_1 + \cdots + m_s e'_s : 0 \leq m_i < n_i, i = 1, 2, \dots, s\}$$

and

$$\mathcal{R}_k = \{\mu_1 e''_1 + \cdots + \mu_s e''_s : 0 \leq \mu_i < n_i, i = 1, 2, \dots, s\},$$

then \mathcal{L}_k and \mathcal{R}_k are sets of all coset representatives of $\mathbb{Z}^s/M^k\mathbb{Z}^s$ and $\mathbb{Z}^s/D^k\mathbb{Z}^s$ respectively. With this particular choice of \mathcal{L}_k , we can order \mathcal{L}_k as in [8], and let us denote $\mathcal{L}_k = \{\ell_1, \dots, \ell_{d^k}\}$.

For each $m, \mu = 1, 2, \dots, r$, consider the $d^k \times d^k$ matrix

$$\Phi_{m\mu} := \left(\langle T_k^{\ell_p} \phi_k^m, T_k^{\ell_q} \tilde{\phi}_k^\mu \rangle \right)_{p,q=1}^{d^k} = \left(\langle \phi_k^m, T_k^{\ell_q - \ell_p} \tilde{\phi}_k^\mu \rangle \right)_{p,q=1}^{d^k},$$

which is a circulant of level s and type (n_1, \dots, n_s) . By [7, Theorem 5.8.4],

$$\Phi_{m\mu} = F^* \left(\sum_{\gamma_1=0}^{n_1-1} \cdots \sum_{\gamma_s=0}^{n_s-1} \langle \phi_k^m, \tilde{\phi}_k^\mu(\cdot - 2\pi M^{-k}(\gamma_1 e'_1 + \cdots + \gamma_s e'_s)) \rangle \right) \Omega^\gamma F. \quad (2.1)$$

Here, $F = F_{n_1} \otimes \cdots \otimes F_{n_s}$ is the Kronecker product of the Fourier matrices F_{n_ν} , $\nu = 1, 2, \dots, s$ (see [7] for the definitions), and $\Omega^\gamma = \Omega_{n_1}^{\gamma_1} \otimes \cdots \otimes \Omega_{n_s}^{\gamma_s}$, where

$$\Omega_{n_\nu} = \text{diag}(1, \omega_{n_\nu}, \omega_{n_\nu}^2, \dots, \omega_{n_\nu}^{n_\nu-1}), \quad \omega_{n_\nu} = \exp(2\pi i/n_\nu).$$

As in [8, Section 3], we deduce that the eigenvalues of $\Phi_{m\mu}$ can be written as

$$\lambda_{m\mu}^j := d^k \sum_{p \in \mathbb{Z}^s} \widehat{\phi}_k^m(j + D^k p) \overline{\widehat{\phi}_k^\mu(j + D^k p)}, \quad j \in \mathcal{R}_k.$$

On the other hand, since

$$\langle v_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle = \sum_{p \in \mathbb{Z}^s} \widehat{\phi}_k^m(j + D^k p) \overline{\widehat{\phi}_k^\mu(j + D^k p)}, \quad j \in \mathcal{R}_k,$$

we have

$$\lambda_{m\mu}^j = d^k \langle v_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle, \quad j \in \mathcal{R}_k. \quad (2.2)$$

Now, if $m \neq \mu$, then $\langle T_k^p \phi_k^m, T_k^q \tilde{\phi}_k^\mu \rangle = 0$ for all $p, q \in \mathcal{L}_k$ amounts to $\Phi_{m\mu} = 0$; and it follows from (2.1) that $\Phi_{m\mu} = 0$ if and only if $\lambda_{m\mu}^j = 0$ for all $j \in \mathcal{R}_k$. By (2.2), this is equivalent to $\langle v_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle = 0$ for all $j \in \mathcal{R}_k$.

If $m = \mu$, then (2.1) implies that

$$\Phi_{mm} = \left(\langle T_k^{\ell_p} \phi_k^m, T_k^{\ell_q} \tilde{\phi}_k^m \rangle \right)_{p,q=1}^{d^k} = I_{d^k}$$

if and only if $\lambda_{mm}^j = 1$ for all $j \in \mathcal{R}_k$ (since F is unitary). By (2.2), this is equivalent to

$$\langle v_{k,j}^m, \tilde{v}_{k,j}^m \rangle = \frac{1}{d^k},$$

which completes the proof of the theorem. \square

3. ALGORITHM FOR MULTIWAVELET CONSTRUCTION

Let $\{\phi_k^m : k \geq 0, m = 1, 2, \dots, r\}$ and $\{\tilde{\phi}_k^m : k \geq 0, m = 1, 2, \dots, r\}$ be two sets of scaling functions that generate a pair of biorthogonal multiresolutions $\{V_k : k \geq 0\}$ and $\{\tilde{V}_k : k \geq 0\}$ of $L^2([0, 2\pi)^s)$ with multiplicity r and dilation matrix M . We shall find functions $\psi_k^m, \tilde{\psi}_k^m \in L^2([0, 2\pi)^s)$, $k \geq 0, m = 1, 2, \dots, r(d-1)$, satisfying the biorthogonality relation (1.4), with the sets W_k and \tilde{W}_k , $k \geq 0$, in (1.5) and (1.6) satisfying (1.7) and (1.8). Our strategy is to construct appropriate polyphase splines in $L^2([0, 2\pi)^s)$ of the form

$$\begin{aligned} u_{k,j}^m(x) &:= \sum_{p \in \mathbb{Z}^s} b_k^m(j + D^k p) e^{i(j + D^k p) \cdot x}, \\ \tilde{u}_{k,j}^m(x) &:= \sum_{p \in \mathbb{Z}^s} \tilde{b}_k^m(j + D^k p) e^{i(j + D^k p) \cdot x}, \end{aligned}$$

for $m = 1, 2, \dots, r(d-1)$, $j \in \mathcal{R}_k$, and then define

$$\psi_k^m := \sum_{j \in \mathcal{R}_k} u_{k,j}^m, \quad \tilde{\psi}_k^m := \sum_{j \in \mathcal{R}_k} \tilde{u}_{k,j}^m,$$

for $m = 1, 2, \dots, r(d-1)$. Here, $b_k^m, \tilde{b}_k^m \in \ell^2(\mathbb{Z}^s)$ are sequences such that

$$\begin{aligned} \left(b_k^1(n), \dots, b_k^{r(d-1)}(n) \right)^T &= \widehat{G}_{k+1}(n) \left(\widehat{\phi}_{k+1}^1(n), \dots, \widehat{\phi}_{k+1}^r(n) \right)^T, \\ \left(\tilde{b}_k^1(n), \dots, \tilde{b}_k^{r(d-1)}(n) \right)^T &= \widetilde{G}_{k+1}(n) \left(\widetilde{\phi}_{k+1}^1(n), \dots, \widetilde{\phi}_{k+1}^r(n) \right)^T, \quad n \in \mathbb{Z}^s, \end{aligned}$$

for some $\widehat{G}_{k+1}, \widetilde{G}_{k+1} \in \mathcal{S}(D^{k+1})^{r(d-1) \times r}$. This is equivalent to

$$\psi_k = \sum_{\ell \in \mathcal{L}_{k+1}} G_{k+1}(\ell) T_{k+1}^\ell \phi_{k+1}, \quad \tilde{\psi}_k = \sum_{\ell \in \mathcal{L}_{k+1}} \widetilde{G}_{k+1}(\ell) T_{k+1}^\ell \tilde{\phi}_{k+1},$$

where $\psi_k := (\psi_k^1, \dots, \psi_k^{r(d-1)})^T$, $\tilde{\psi}_k := (\tilde{\psi}_k^1, \dots, \tilde{\psi}_k^{r(d-1)})^T$. Note that the sets W_k and \widetilde{W}_k in (1.5) and (1.6) can be expressed as

$$W_k = \langle \{u_{k,j}^m : m = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\} \rangle$$

and

$$\widetilde{W}_k = \langle \{\tilde{u}_{k,j}^m : m = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\} \rangle.$$

(See [8] for details on the above general approach of constructing multiwavelets via polyphase splines.)

The desired polyphase splines, and hence multiwavelets, will be obtained by constructing matrices $\widehat{G}_{k+1}, \widetilde{G}_{k+1} \in \mathcal{S}(D^{k+1})^{r(d-1) \times r}$ so that (1.4), (1.7) and (1.8) are satisfied. For each $k \geq 0$, $j \in \mathcal{R}_k$, we define the following matrices:

$$\begin{aligned} M_k(j) &:= \left(\langle v_{k,j}^m, v_{k,j}^\mu \rangle \right)_{m,\mu=1}^r, & \widetilde{M}_k(j) &:= \left(\langle \tilde{v}_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle \right)_{m,\mu=1}^r, \\ N_k(j) &:= \left(\langle u_{k,j}^m, u_{k,j}^\mu \rangle \right)_{m,\mu=1}^{r(d-1)}, & \widetilde{N}_k(j) &:= \left(\langle \tilde{u}_{k,j}^m, \tilde{u}_{k,j}^\mu \rangle \right)_{m,\mu=1}^{r(d-1)}, \\ \mathcal{M}_k(j) &:= \left(\langle v_{k,j}^m, \tilde{v}_{k,j}^\mu \rangle \right)_{m,\mu=1}^r, & \mathcal{N}_k(j) &:= \left(\langle u_{k,j}^m, \tilde{u}_{k,j}^\mu \rangle \right)_{m,\mu=1}^{r(d-1)}. \end{aligned}$$

By the arguments in [8], for $j \in \mathcal{R}_k$, we have

$$M_k(j) = \sum_{\ell \in \mathcal{R}_1} \widehat{H}_{k+1}(j + D^k \ell) M_{k+1}(j + D^k \ell) \widehat{H}_{k+1}(j + D^k \ell)^*, \quad (3.1)$$

$$N_k(j) = \sum_{\ell \in \mathcal{R}_1} \widehat{G}_{k+1}(j + D^k \ell) M_{k+1}(j + D^k \ell) \widehat{G}_{k+1}(j + D^k \ell)^*, \quad (3.2)$$

as well as the corresponding identities for $\widetilde{M}_k(j)$ and $\widetilde{N}_k(j)$. Since $\{V_k : k \geq 0\}$ and $\{\widetilde{V}_k : k \geq 0\}$ satisfy MR1, by [8, Propositions 3.1 and 3.2], we obtain

$$\det(M_k(j)) > 0, \quad \det(\widetilde{M}_k(j)) > 0, \quad k \geq 0, \quad j \in \mathcal{R}_k. \quad (3.3)$$

Analogous to (3.1) and (3.2), we can derive the following identities:

$$\mathcal{M}_k(j) = \sum_{\ell \in \mathcal{R}_1} \widehat{H}_{k+1}(j + D^k \ell) \mathcal{M}_{k+1}(j + D^k \ell) \widehat{H}_{k+1}(j + D^k \ell)^*, \quad (3.4)$$

$$\mathcal{N}_k(j) = \sum_{\ell \in \mathcal{R}_1} \widehat{G}_{k+1}(j + D^k \ell) \mathcal{M}_{k+1}(j + D^k \ell) \widehat{G}_{k+1}(j + D^k \ell)^*. \quad (3.5)$$

By Theorem 2.1, (1.2) is equivalent to

$$\mathcal{M}_k(j) = \frac{1}{d^k} I_r, \quad k \geq 0, \quad j \in \mathcal{R}_k. \quad (3.6)$$

Therefore (3.4) implies that

$$\sum_{\ell \in \mathcal{R}_1} \widehat{H}_{k+1}(j + D^k \ell) \widehat{H}_{k+1}(j + D^k \ell)^* = d I_r. \quad (3.7)$$

By Theorem 2.1 again, we see that (1.4) is equivalent to

$$\mathcal{N}_k(j) = \frac{1}{d^k} I_{r(d-1)}, \quad k \geq 0, \quad j \in \mathcal{R}_k,$$

which, by (3.5) and (3.6), amounts to

$$\sum_{\ell \in \mathcal{R}_1} \widehat{G}_{k+1}(j + D^k \ell) \widehat{G}_{k+1}(j + D^k \ell)^* = d I_{r(d-1)}. \quad (3.8)$$

As for the direct sums in (1.7), note that it suffices to establish the linear independence of the sets $\{v_{k,j}^m, u_{k,j}^\mu : m = 1, 2, \dots, r, \mu = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\}$ and $\{\widetilde{v}_{k,j}^m, \widetilde{u}_{k,j}^\mu : m = 1, 2, \dots, r, \mu = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\}$. Writing

$$P_k(j) := (\langle v_{k,j}^m, u_{k,j}^\mu \rangle)_{m=1, \mu=1}^{r, r(d-1)}, \quad \widetilde{P}_k(j) := (\langle \widetilde{v}_{k,j}^m, \widetilde{u}_{k,j}^\mu \rangle)_{m=1, \mu=1}^{r, r(d-1)}, \quad k \geq 0, \quad j \in \mathcal{R}_k, \quad (3.9)$$

we claim that the desired linear independence is equivalent to

$$\det \left(\begin{array}{c|c} M_k(j) & P_k(j) \\ \hline P_k(j)^* & N_k(j) \end{array} \right) > 0, \quad \det \left(\begin{array}{c|c} \widetilde{M}_k(j) & \widetilde{P}_k(j) \\ \hline \widetilde{P}_k(j)^* & \widetilde{N}_k(j) \end{array} \right) > 0, \quad k \geq 0, \quad j \in \mathcal{R}_k. \quad (3.10)$$

Indeed, observe that

$$\left(\begin{array}{c|c} M_k(j) & P_k(j) \\ \hline P_k(j)^* & N_k(j) \end{array} \right)$$

is just the Gram matrix of $\mathcal{B}_j := \{v_{k,j}^m, u_{k,j}^\mu : m = 1, 2, \dots, r, \mu = 1, 2, \dots, r(d-1)\}$. Hence, (3.10) implies that \mathcal{B}_j is linearly independent for each fixed $j \in \mathcal{R}_k$. As $\langle v_{k,j}^m, u_{k,\ell}^\mu \rangle = \langle v_{k,j}^m, v_{k,\ell}^{m'} \rangle = \langle u_{k,j}^\mu, u_{k,\ell}^{\mu'} \rangle = 0$ for all $m, m' = 1, 2, \dots, r, \mu, \mu' = 1, 2, \dots, r(d-1)$ if $j \neq \ell$, we see that \mathcal{B}_j is orthogonal to \mathcal{B}_ℓ if $j \neq \ell, j, \ell \in \mathcal{R}_k$. It follows that

$$\bigcup_{j \in \mathcal{R}_k} \mathcal{B}_j = \{v_{k,j}^m, u_{k,j}^\mu : m = 1, 2, \dots, r, \mu = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\}$$

is a linearly independent set. Similarly, we obtain the result for the other set of polyphase splines.

As a consequence of the linear independence condition characterized by (3.10), each of the collections $\{u_{k,j}^m : m = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\}$ and $\{\widetilde{u}_{k,j}^m : m = 1, 2, \dots, r(d-1), j \in \mathcal{R}_k\}$ is linearly independent. In this case, the orthogonality of W_k and \widetilde{V}_k translates into

$$\langle u_{k,j}^m, \widetilde{v}_{k,\ell}^\mu \rangle = 0, \quad m = 1, 2, \dots, r(d-1), \quad \mu = 1, 2, \dots, r, \quad j, \ell \in \mathcal{R}_k,$$

which is equivalent to

$$\langle u_{k,j}^m, \widetilde{v}_{k,j}^\mu \rangle = 0, \quad m = 1, 2, \dots, r(d-1), \quad \mu = 1, 2, \dots, r, \quad j \in \mathcal{R}_k. \quad (3.11)$$

Proceeding as in [8], we deduce that (3.11) can be written as

$$\sum_{\ell \in \mathcal{R}_1} \widehat{G}_{k+1}(j + D^k \ell) \mathcal{M}_{k+1}(j + D^k \ell) \widehat{H}_{k+1}(j + D^k \ell)^* = 0_{r(d-1) \times r},$$

which, by (3.6), is equivalent to

$$\sum_{\ell \in \mathcal{R}_1} \widehat{G}_{k+1}(j + D^k \ell) \widetilde{\widehat{H}}_{k+1}(j + D^k \ell)^* = 0_{r(d-1) \times r}. \quad (3.12)$$

Similarly, \widetilde{W}_k being orthogonal to V_k amounts to

$$\sum_{\ell \in \mathcal{R}_1} \widetilde{\widehat{G}}_{k+1}(j + D^k \ell) \widehat{H}_{k+1}(j + D^k \ell)^* = 0_{r(d-1) \times r}. \quad (3.13)$$

Hence, the multiwavelet construction problem can now be formulated as a problem of constructing matrices $\widehat{G}_{k+1}, \widetilde{\widehat{G}}_{k+1} \in \mathcal{S}(D^{k+1})^{r(d-1) \times r}$ satisfying (3.8), (3.10), (3.12) and (3.13), under the assumptions of (3.1), (3.3) and (3.7). Our main theorem, Theorem 3.1, shows that such matrices can always be constructed.

Theorem 3.1. *Suppose that $\phi_k^m, \widetilde{\phi}_k^m \in L^2([0, 2\pi]^s)$, $k \geq 0$, $m = 1, 2, \dots, r$, generate a pair of biorthogonal multiresolutions of $L^2([0, 2\pi]^s)$ with multiplicity r and dilation matrix M , so that (3.1), (3.3) and (3.7) hold. Then there exist $\widehat{G}_{k+1}, \widetilde{\widehat{G}}_{k+1} \in \mathcal{S}(D^{k+1})^{r(d-1) \times r}$ such that (3.8), (3.10), (3.12) and (3.13) are satisfied.*

Before we prove the theorem, let us establish a useful technical lemma.

Lemma 3.1. *Suppose that A and \widetilde{A} are $r \times n$ complex matrices, where $r < n$, satisfying*

$$A\widetilde{A}^* = I_r. \quad (3.14)$$

Then there exist $n \times n$ complex matrices C and \widetilde{C} of the form

$$C = \begin{pmatrix} A \\ B \end{pmatrix}, \quad \widetilde{C} = \begin{pmatrix} \widetilde{A} \\ \widetilde{B} \end{pmatrix},$$

for some $(n-r) \times n$ complex matrices B and \widetilde{B} such that

$$C\widetilde{C}^* = I_n.$$

Proof. Let v_1, \dots, v_r be the rows of A . Since $A\widetilde{A}^* = I_r$, it follows that the rank of A is r . Thus there exist $(n-r)$ linearly independent vectors $\widetilde{w}_1, \dots, \widetilde{w}_{n-r}$ which form

a basis of the orthogonal complement of the linear span of $\{v_1, \dots, v_r\}$ in \mathbb{C}^n . Define an $n \times n$ complex matrix \tilde{C} by

$$\tilde{C} = \begin{pmatrix} \tilde{A} \\ \tilde{B} \end{pmatrix},$$

where \tilde{B} is the $(n-r) \times n$ matrix formed by the vectors $\tilde{w}_1, \dots, \tilde{w}_{n-r}$.

We claim that the matrix \tilde{C} is invertible. Indeed, if

$$\alpha_1 \tilde{v}_1 + \dots + \alpha_r \tilde{v}_r + \beta_1 \tilde{w}_1 + \dots + \beta_{n-r} \tilde{w}_{n-r} = 0_{1 \times n}$$

for some $\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_{n-r} \in \mathbb{C}$, then taking inner product with v_1, \dots, v_r respectively yields $\alpha_1 = \dots = \alpha_r = 0$. Since $\{\tilde{w}_1, \dots, \tilde{w}_{n-r}\}$ is linearly independent, it follows that $\beta_1 = \dots = \beta_{n-r} = 0$. Thus $\{\tilde{v}_1, \dots, \tilde{v}_r, \tilde{w}_1, \dots, \tilde{w}_{n-r}\}$ is linearly independent.

Now, set $C_0 := (\tilde{C}^*)^{-1}$. Writing $C_0 = \begin{pmatrix} A_0 \\ B \end{pmatrix}$ where A_0 and B are $r \times n$ and $(n-r) \times n$ matrices respectively, the relation $C_0 \tilde{C}^* = I_n$ implies that

$$B \tilde{A}^* = 0_{(n-r) \times r}, \quad B \tilde{B}^* = I_{n-r}. \quad (3.15)$$

Define $C := \begin{pmatrix} A \\ B \end{pmatrix}$. By (3.14), (3.15), and the fact that the rows of \tilde{B} form a basis of the orthogonal complement of the linear span of the rows of A in \mathbb{C}^n , we have $C \tilde{C}^* = I_n$. (This also shows that $C = (\tilde{C}^*)^{-1} = C_0$.) Hence, the proof of the lemma is complete. \square

Proof of Theorem 3.1. Fix $k \geq 0$, $j \in \mathcal{R}_k$, and let $\ell_1, \ell_2, \dots, \ell_d$ denote all the elements of \mathcal{R}_1 . Set

$$\begin{aligned} A_k(j) &:= \frac{1}{\sqrt{d}} (\widehat{H}_{k+1}(j + D^k \ell_1) \mid \dots \mid \widehat{H}_{k+1}(j + D^k \ell_d)), \\ \tilde{A}_k(j) &:= \frac{1}{\sqrt{d}} (\widehat{\tilde{H}}_{k+1}(j + D^k \ell_1) \mid \dots \mid \widehat{\tilde{H}}_{k+1}(j + D^k \ell_d)). \end{aligned} \quad (3.16)$$

Then (3.7) is equivalent to

$$A_k(j) \tilde{A}_k(j)^* = I_r.$$

By Lemma 3.1, there exist $rd \times rd$ matrices $C_k(j)$ and $\tilde{C}_k(j)$ of the form

$$C_k(j) = \begin{pmatrix} A_k(j) \\ B_k(j) \end{pmatrix}, \quad \tilde{C}_k(j) = \begin{pmatrix} \tilde{A}_k(j) \\ \tilde{B}_k(j) \end{pmatrix}, \quad (3.17)$$

for some $r(d-1) \times rd$ matrices $B_k(j)$ and $\tilde{B}_k(j)$ such that

$$C_k(j)\tilde{C}_k(j)^* = I_{rd}. \quad (3.18)$$

Now, define $r(d-1) \times r$ matrices $\hat{G}_{k+1}(j + D^k \ell)$, $\tilde{\hat{G}}_{k+1}(j + D^k \ell)$, $\ell \in \mathcal{R}_1$, by

$$\begin{aligned} B_k(j) &:= \frac{1}{\sqrt{d}}(\hat{G}_{k+1}(j + D^k \ell_1) \mid \cdots \mid \hat{G}_{k+1}(j + D^k \ell_d)), \\ \tilde{B}_k(j) &:= \frac{1}{\sqrt{d}}(\tilde{\hat{G}}_{k+1}(j + D^k \ell_1) \mid \cdots \mid \tilde{\hat{G}}_{k+1}(j + D^k \ell_d)). \end{aligned} \quad (3.19)$$

By (3.17) and (3.18), we have

$$\left(\begin{array}{c|c} A_k(j)\tilde{A}_k(j)^* & A_k(j)\tilde{B}_k(j)^* \\ \hline B_k(j)\tilde{A}_k(j)^* & B_k(j)\tilde{B}_k(j)^* \end{array} \right) = \left(\begin{array}{c|c} I_r & 0_{r \times r(d-1)} \\ \hline 0_{r(d-1) \times r} & I_{r(d-1)} \end{array} \right),$$

and it follows from (3.16) and (3.19) that the conditions (3.8), (3.12) and (3.13) are satisfied.

It remains to show that (3.10) holds. Note that the matrix $P_k(j)$ defined in (3.9) can be written as

$$P_k(j) = \sum_{\ell \in \mathcal{R}_1} \hat{H}_{k+1}(j + D^k \ell) M_{k+1}(j + D^k \ell) \tilde{\hat{G}}_{k+1}(j + D^k \ell)^*.$$

Then by (3.1) and (3.2), we see that

$$C_k(j) \text{diag}(M_{k+1}(j + D^k \ell_1), \dots, M_{k+1}(j + D^k \ell_d)) C_k(j)^* = \frac{1}{d} \left(\begin{array}{c|c} M_k(j) & P_k(j) \\ \hline P_k(j)^* & N_k(j) \end{array} \right).$$

Since $C_k(j)$ is invertible and (3.3) holds, the determinant of the product of matrices in the above identity is a positive real number. Consequently, we have

$$\det \left(\begin{array}{c|c} M_k(j) & P_k(j) \\ \hline P_k(j)^* & N_k(j) \end{array} \right) > 0.$$

A similar argument shows that

$$\det \left(\begin{array}{c|c} \widetilde{M}_k(j) & \widetilde{P}_k(j) \\ \hline \widetilde{P}_k(j)^* & \widetilde{N}_k(j) \end{array} \right) > 0,$$

and this completes the proof of the theorem. \square

The proof of Theorem 3.1 is constructive, and it gives an algorithm for the construction of biorthogonal multiwavelets from a pair of biorthogonal multiresolutions.

Algorithm 3.1. For $k \geq 0$, suppose that $\widehat{H}_{k+1}, \widetilde{H}_{k+1} \in \mathcal{S}(D^{k+1})^{r \times r}$ are the matrices in the periodic matrix refinement equations (1.3).

Step 1: Fix $j \in \mathcal{R}_k$ and form the $r \times rd$ matrices $A_k(j)$ and $\widetilde{A}_k(j)$ as in (3.16).

Step 2: Find a $(rd - r) \times rd$ matrix $\widetilde{B}_k(j)$ whose rows formed a basis of the orthogonal complement of the linear span of the rows of $A_k(j)$ in \mathbb{C}^{rd} .

Step 3: Set $B_k(j)$ to be the $(rd - r) \times rd$ matrix formed by the last $(rd - r)$ rows of the $rd \times rd$ matrix $(\widetilde{A}_k(j)^* | \widetilde{B}_k(j)^*)^{-1}$.

Step 4: Define $r(d - 1) \times r$ matrices $\widehat{G}_{k+1}(j + D^k \ell), \widetilde{G}_{k+1}(j + D^k \ell), \ell \in \mathcal{R}_1$, by (3.19).

Step 5: Repeat Steps 1–4 with all the elements j in \mathcal{R}_k .

Step 6: Write $\psi_k = (\psi_k^1, \dots, \psi_k^{r(d-1)})^T$ and $\widetilde{\psi}_k = (\widetilde{\psi}_k^1, \dots, \widetilde{\psi}_k^{r(d-1)})^T$, and set

$$\begin{aligned} \psi_k(x) &:= \sum_{n \in \mathbb{Z}^s} \widehat{G}_{k+1}(n) \left(\widehat{\phi}_{k+1}^1(n), \dots, \widehat{\phi}_{k+1}^r(n) \right)^T e^{in \cdot x}, \\ \widetilde{\psi}_k(x) &:= \sum_{n \in \mathbb{Z}^s} \widetilde{G}_{k+1}(n) \left(\widetilde{\phi}_{k+1}^1(n), \dots, \widetilde{\phi}_{k+1}^r(n) \right)^T e^{in \cdot x}, \end{aligned}$$

to obtain the biorthogonal multiwavelets in $L^2([0, 2\pi]^s)$.

In the following two examples on the special case of $r = 1$, different values of d are considered.

Example 3.1. Let $d = 2$. Then

(a) $s = 1, M = 2$,

$$(b) \ s = 2, \ M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \text{ or } \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \text{ or } \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix},$$

are some examples of the cases included under this setting. Without loss of generality, we may assume that $\mathcal{L}_1 = \{0, q_2\}$ and $\mathcal{R}_1 = \{0, \ell_2\}$.

For $k \geq 0$, $j \in \mathcal{R}_k$, define

$$\begin{aligned} \widehat{G}_{k+1}(j) &= -\frac{1}{\sqrt{2}} e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j + D^k \ell_2)}, \\ \widehat{G}_{k+1}(j + D^k \ell_2) &= \frac{1}{\sqrt{2}} e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j)}. \end{aligned} \quad (3.20)$$

As shown independently in [3] and [10], $e^{-i\ell_2 \cdot 2\pi M^{-1} q_2} = -1$. This implies that the two equations in (3.20) can be combined together as

$$\widehat{G}_{k+1}(j) = -\frac{1}{\sqrt{2}} e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j + D^k \ell_2)}, \quad j \in \mathcal{R}_{k+1}. \quad (3.21)$$

Now, for $k \geq 0$, $j \in \mathcal{R}_k$, the matrix $\widetilde{C}_k(j)$ in (3.17) is given by

$$\widetilde{C}_k(j) = \frac{1}{\sqrt{2}} \begin{pmatrix} \widehat{H}_{k+1}(j) & \widehat{H}_{k+1}(j + D^k \ell_2) \\ -e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j + D^k \ell_2)} & e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j)} \end{pmatrix}.$$

Then the matrix $C_k(j) = (\widetilde{C}_k(j)^*)^{-1}$ in (3.17) yields

$$\widehat{G}_{k+1}(j) = -\frac{1}{\sqrt{2}} e^{-ij \cdot 2\pi M^{-(k+1)} q_2} \overline{\widehat{H}_{k+1}(j + D^k \ell_2)}, \quad j \in \mathcal{R}_{k+1}. \quad (3.22)$$

Note that (3.21) and (3.22) are equivalent to

$$\widetilde{G}_{k+1}(\ell) = e^{-i\ell_2 \cdot 2\pi M^{-1} \ell} \overline{\widehat{H}_{k+1}(q_2 - \ell)}, \quad G_{k+1}(\ell) = e^{-i\ell_2 \cdot 2\pi M^{-1} \ell} \overline{\widetilde{H}_{k+1}(q_2 - \ell)},$$

for all $\ell \in \mathcal{L}_{k+1}$. In the special case of $s = 1$ and $M = 2$, this becomes

$$\widetilde{G}_{k+1}(\ell) = (-1)^\ell \overline{\widehat{H}_{k+1}(1 - \ell)}, \quad G_{k+1}(\ell) = (-1)^\ell \overline{\widetilde{H}_{k+1}(1 - \ell)},$$

for all $\ell = 0, 1, \dots, 2^{k+1} - 1$. These expressions are periodic analogues of the familiar alternating flip formulas for the construction of wavelets in $L^2(\mathbb{R})$. \square

Example 3.2. Let $d > 2$, and fix $k \geq 0$, $j \in \mathcal{R}_k$. Without loss of generality, suppose that $\ell_1, \ell_2, \dots, \ell_d$ are all the elements of \mathcal{R}_1 with

$$|\widehat{H}_{k+1}(j + D^k \ell_1)| = \max_{\ell \in \mathcal{R}_1} |\widehat{H}_{k+1}(j + D^k \ell)|.$$

Define the $(d-1) \times 1$ matrices

$$\widehat{\widetilde{G}}_{k+1}(j + D^k \ell) := \left(\widehat{\widetilde{G}}_{k+1}^1(j + D^k \ell), \dots, \widehat{\widetilde{G}}_{k+1}^{d-1}(j + D^k \ell) \right)^T, \quad \ell \in \mathcal{R}_1,$$

by

$$\widehat{\widetilde{G}}_{k+1}^m(j + D^k \ell_\nu) = \begin{cases} -\frac{1}{\sqrt{d}} \overline{\widehat{H}_{k+1}(j + D^k \ell_{m+1})}, & \text{if } \nu = 1, \\ \frac{1}{\sqrt{d}} \overline{\widehat{H}_{k+1}(j + D^k \ell_1)}, & \text{if } \nu = m + 1, \\ 0, & \text{otherwise,} \end{cases}$$

where $m = 1, 2, \dots, d-1$, $\nu = 1, 2, \dots, d$. Then the matrix $\widetilde{C}_k(j)$ in (3.17) is given by

$$\widetilde{C}_k(j) = \frac{1}{\sqrt{d}} \times \begin{pmatrix} \overline{\widehat{H}_{k+1}(j + D^k \ell_1)} & \overline{\widehat{H}_{k+1}(j + D^k \ell_2)} & \overline{\widehat{H}_{k+1}(j + D^k \ell_3)} & \cdots & \overline{\widehat{H}_{k+1}(j + D^k \ell_d)} \\ -\overline{\widehat{H}_{k+1}(j + D^k \ell_2)} & \overline{\widehat{H}_{k+1}(j + D^k \ell_1)} & 0 & \cdots & 0 \\ -\overline{\widehat{H}_{k+1}(j + D^k \ell_3)} & 0 & \overline{\widehat{H}_{k+1}(j + D^k \ell_1)} & \cdots & 0 \\ \vdots & \vdots & \cdots & \ddots & \vdots \\ -\overline{\widehat{H}_{k+1}(j + D^k \ell_d)} & 0 & \cdots & 0 & \overline{\widehat{H}_{k+1}(j + D^k \ell_1)} \end{pmatrix},$$

and a direct calculation yields

$$\det(\widetilde{C}_k(j)) = \left(\frac{1}{\sqrt{d}} \right)^d \overline{\widehat{H}_{k+1}(j + D^k \ell_1)}^{d-2} \sum_{\nu=1}^d \widehat{H}_{k+1}(j + D^k \ell_\nu) \overline{\widehat{H}_{k+1}(j + D^k \ell_\nu)}.$$

Consequently, it follows from (3.7) that

$$\det(\widetilde{C}_k(j)) = d^{1-\frac{d}{2}} \overline{\widehat{H}_{k+1}(j + D^k \ell_1)}^{d-2} \neq 0,$$

since $|\widehat{H}_{k+1}(j + D^k \ell_1)| = \max_{\ell \in \mathcal{R}_1} |\widehat{H}_{k+1}(j + D^k \ell)| > 0$.

The $(d-1) \times 1$ matrices $\widehat{G}_{k+1}(j + D^k \ell)$, $\ell \in \mathcal{R}_1$, are obtained from the matrix $C_k(j)$ in (3.17) which is given by

$$C_k(j) = (\widetilde{C}_k(j)^*)^{-1} = \frac{1}{d^{1-\frac{d}{2}} \overline{\widehat{H}_{k+1}(j + D^k \ell_1)}^{d-2}} \text{adj}(\widetilde{C}_k(j)^*),$$

where $\text{adj}(\widetilde{C}_k(j)^*)$ denotes the transpose of the matrix of cofactors of $\widetilde{C}_k(j)^*$. \square

4. BIORTHOGONAL RIESZ BASES

For $k \geq 0$, suppose that $\{T_k^\ell \psi_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\}$ and $\{T_k^\ell \widetilde{\psi}_k^m : m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\}$ form a pair of biorthogonal bases of W_k and \widetilde{W}_k respectively. Since $L^2([0, 2\pi]^s)$ has the direct sum decompositions

$$L^2([0, 2\pi]^s) = V_0 \oplus W_0 \oplus W_1 \oplus \dots = \widetilde{V}_0 \oplus \widetilde{W}_0 \oplus \widetilde{W}_1 \oplus \dots,$$

it is natural to ask whether the two collections of functions

$$\{\phi_0^m : m = 1, 2, \dots, r\} \cup \{T_k^\ell \psi_k^m : k \geq 0, m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\} \quad (4.1)$$

and

$$\{\widetilde{\phi}_0^m : m = 1, 2, \dots, r\} \cup \{T_k^\ell \widetilde{\psi}_k^m : k \geq 0, m = 1, 2, \dots, r(d-1), \ell \in \mathcal{L}_k\} \quad (4.2)$$

form a pair of biorthogonal Riesz bases of $L^2([0, 2\pi]^s)$. In this connection, the following theorem, which is a multidimensional periodic analogue of results in [3], [5], [6], on a general sequence of functions ψ_k^m , $k \geq k_0$, $m = 1, 2, \dots, r(d-1)$, is useful.

Theorem 4.1. *Suppose that there exist positive constants C_1 and C_2 such that*

$$\sum_{k=k_0}^{\infty} \sum_{m=1}^{r(d-1)} |\sqrt{d^k} \widehat{\psi}_k^m(n)|^\epsilon \leq C_1, \quad n \in \mathbb{Z}^s, \quad (4.3)$$

and

$$\sum_{p \in \mathbb{Z}^s} |\sqrt{d^k} \widehat{\psi}_k^m(j + D^k p)|^{2-\epsilon} \leq C_2, \quad k \geq k_0, \quad m = 1, 2, \dots, r(d-1), \quad j \in \mathcal{R}_k, \quad (4.4)$$

for some $k_0 \geq 0$ and $\epsilon > 0$. Then there exists a positive constant B_0 such that for every $f \in L^2([0, 2\pi]^s)$,

$$\sum_{k=k_0}^{\infty} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 \leq B_0 \|f\|^2. \quad (4.5)$$

Proof. First, fix $k \geq k_0$. By the Parseval's identity,

$$\begin{aligned} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 &= \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} \left| \sum_{n \in \mathbb{Z}^s} \widehat{f}(n) \overline{\widehat{\psi}_k^m(n)} e^{in \cdot (2\pi M^{-k} \ell)} \right|^2 \\ &= \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} \left| \sum_{j \in \mathcal{R}_k} \sum_{p \in \mathbb{Z}^s} \widehat{f}(j + D^k p) \overline{\widehat{\psi}_k^m(j + D^k p)} e^{ij \cdot (2\pi M^{-k} \ell)} \right|^2 \end{aligned} \quad (4.6)$$

Let

$$\widehat{\alpha}_k^m(j) := \sum_{p \in \mathbb{Z}^s} \widehat{f}(j + D^k p) \overline{\widehat{\psi}_k^m(j + D^k p)}, \quad j \in \mathcal{R}_k.$$

Then $\widehat{\alpha}_k^m \in \mathcal{S}(D^k)$, and its inverse finite Fourier transform α_k^m is a sequence in $\mathcal{S}(M^k)$. Furthermore, $\widehat{\alpha}_k^m$ and α_k^m satisfy the relation (1.1). Consequently, it follows from (4.6) that

$$\begin{aligned} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 &= \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} \left| \sum_{j \in \mathcal{R}_k} \widehat{\alpha}_k^m(j) e^{ij \cdot (2\pi M^{-k} \ell)} \right|^2 \\ &= \sum_{m=1}^{r(d-1)} |\mathcal{R}_k|^2 \sum_{\ell \in \mathcal{L}_k} |\alpha_k^m(\ell)|^2 \\ &= \sum_{m=1}^{r(d-1)} |\mathcal{R}_k| \sum_{j \in \mathcal{R}_k} |\widehat{\alpha}_k^m(j)|^2. \end{aligned}$$

This implies that for $\epsilon > 0$,

$$\begin{aligned} &\sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 \\ &\leq d^k \sum_{m=1}^{r(d-1)} \sum_{j \in \mathcal{R}_k} \left(\sum_{p \in \mathbb{Z}^s} |\widehat{f}(j + D^k p)| \cdot |\widehat{\psi}_k^m(j + D^k p)|^{\frac{\epsilon}{2}} \cdot |\widehat{\psi}_k^m(j + D^k p)|^{1-\frac{\epsilon}{2}} \right)^2 \\ &\leq d^k \sum_{m=1}^{r(d-1)} \sum_{j \in \mathcal{R}_k} \left(\sum_{p \in \mathbb{Z}^s} |\widehat{f}(j + D^k p)|^2 |\widehat{\psi}_k^m(j + D^k p)|^\epsilon \right) \cdot \left(\sum_{p \in \mathbb{Z}^s} |\widehat{\psi}_k^m(j + D^k p)|^{2-\epsilon} \right) \\ &\leq C_2 \sum_{m=1}^{r(d-1)} \sum_{j \in \mathcal{R}_k} \sum_{p \in \mathbb{Z}^s} |\widehat{f}(j + D^k p)|^2 |\sqrt{d^k} \widehat{\psi}_k^m(j + D^k p)|^\epsilon, \end{aligned}$$

where the assumption (4.4) is used in the last inequality.

Hence,

$$\begin{aligned} \sum_{k=k_0}^{\infty} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 &\leq C_2 \sum_{k=k_0}^{\infty} \sum_{m=1}^{r(d-1)} \sum_{n \in \mathbb{Z}^s} |\widehat{f}(n)|^2 |\sqrt{d^k} \widehat{\psi}_k^m(n)|^\epsilon, \\ &= C_2 \sum_{n \in \mathbb{Z}^s} |\widehat{f}(n)|^2 \sum_{k=k_0}^{\infty} \sum_{m=1}^{r(d-1)} |\sqrt{d^k} \widehat{\psi}_k^m(n)|^\epsilon. \end{aligned}$$

Then (4.5) follows from the assumption (4.3). \square

Theorem 4.2. *For each of the collections of functions (4.1) and (4.2), suppose that there exist positive constants C_1 and C_2 such that (4.3) and (4.4) are satisfied for some $k_0 \geq 0$ and $\epsilon > 0$. Then (4.1) and (4.2) form a pair of biorthogonal Riesz bases of $L^2([0, 2\pi]^s)$.*

Proof. By a standard argument (see [5] or [14, pp. 32–36]), it suffices to establish the existence of a positive constant B such that the inequality

$$\sum_{m=1}^d |\langle f, \phi_0^m \rangle|^2 + \sum_{k=0}^{\infty} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 \leq B \|f\|^2, \quad f \in L^2([0, 2\pi]^s),$$

holds for both collections (4.1) and (4.2). Indeed, it follows from the Cauchy-Schwarz's inequality and Theorem 4.1 that for the collection (4.1),

$$\begin{aligned} &\sum_{m=1}^d |\langle f, \phi_0^m \rangle|^2 + \sum_{k=0}^{\infty} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} |\langle f, T_k^\ell \psi_k^m \rangle|^2 \\ &\leq \left(\sum_{m=1}^d \|\phi_0^m\|^2 + \sum_{k=0}^{k_0-1} \sum_{m=1}^{r(d-1)} \sum_{\ell \in \mathcal{L}_k} \|T_k^\ell \psi_k^m\|^2 + B_0 \right) \|f\|^2, \end{aligned}$$

where the constant B_0 is as in (4.5). Similarly, the collection (4.2) satisfies another such inequality. Thus the proof of the theorem is complete. \square

For the special case of $r = 1$ and $d = 2$ on the space $L^2(\mathbb{R}^s)$, the compactly supported biorthogonal wavelets ψ and $\widetilde{\psi}$ in [5], [6] satisfy the conditions

$$\sup_{\omega \in \mathbb{R}^s} \sum_{k=-\infty}^{\infty} |\widehat{\psi}(D^k \omega)|^\epsilon < \infty, \quad \sup_{\omega \in \mathbb{R}^s} \sum_{k=-\infty}^{\infty} |\widehat{\widetilde{\psi}}(D^k \omega)|^{\widetilde{\epsilon}} < \infty, \quad (4.7)$$

and

$$\sup_{\omega \in \mathbb{R}^s} \sum_{p \in \mathbb{Z}^s} |\widehat{\psi}(\omega + 2\pi p)|^{2-\epsilon} < \infty, \quad \sup_{\omega \in \mathbb{R}^s} \sum_{p \in \mathbb{Z}^s} |\widetilde{\psi}(\omega + 2\pi p)|^{2-\tilde{\epsilon}} < \infty, \quad (4.8)$$

where ϵ and $\tilde{\epsilon}$ are positive constants. For $k \geq 0$, let ψ_k and $\widetilde{\psi}_k$ be biorthogonal wavelets in $L^2([0, 2\pi)^s)$ obtained from periodizing ψ and $\widetilde{\psi}$ with the formulas:

$$\psi_k(x) = \sqrt{d^k} \sum_{p \in \mathbb{Z}^s} \psi \left(\frac{M^k}{2\pi} x + M^k p \right), \quad \widetilde{\psi}_k(x) = \sqrt{d^k} \sum_{p \in \mathbb{Z}^s} \widetilde{\psi} \left(\frac{M^k}{2\pi} x + M^k p \right).$$

Then it follows from the conditions (4.7) and (4.8) that ψ_k and $\widetilde{\psi}_k$ satisfy (4.3) and (4.4) for some k_0 . On a related note, suppose that ϕ_k and $\widetilde{\phi}_k$, $k \geq 0$, are biorthogonal scaling functions in $L^2([0, 2\pi)^s)$ obtained by periodizing the compactly supported biorthogonal scaling functions of $L^2(\mathbb{R}^s)$ in [5], [6]. Then the matrix extension procedure in Example 3.1 also yields the same biorthogonal wavelets ψ_k and $\widetilde{\psi}_k$, $k \geq 0$, which satisfy (4.3) and (4.4).

There are still many open problems in the topic of multidimensional biorthogonal periodic multiwavelets. For instance, for the space $L^2(\mathbb{R}^s)$, the study of biorthogonal scaling functions and multiwavelets can be formulated in terms of transition operators and cascade algorithms. It would be interesting to derive such an approach for the space $L^2([0, 2\pi)^s)$.

Acknowledgement. We would like to thank Professor S. L. Lee of the National University of Singapore and Professor Rong-Qing Jia of the University of Alberta for helpful suggestions and insightful discussions.

REFERENCES

- [1] H. L. Chen, Wavelets from trigonometric spline approach, *J. Approx. Theory and Appl.* **12** (1996), 99–110.
- [2] H. L. Chen, Construction of orthonormal wavelets in the periodic case, *Chinese Sci. Bull.* **14** (1996), 552–554.
- [3] C. K. Chui and C. Li, A general framework of multivariate wavelets with duals, *Appl. Comput. Harmonic Anal.* **1** (1994), 368–390.
- [4] C. K. Chui and N. H. Mhaskar, On trigonometric wavelets, *Constr. Approx.* **9** (1993), 167–190.

- [5] A. Cohen and I. Daubechies, A stability criterion for biorthogonal wavelet bases and their related subband coding scheme, *Duke Math. J.* **68** (1992), 313–335.
- [6] A. Cohen and I. Daubechies, Non-separable bidimensional wavelet bases, *Rev. Mat. Iberoamericana* **9** (1993), 51–137.
- [7] P. J. Davis, *Circulant Matrices* (Wiley, 1979).
- [8] S. S. Goh, S. L. Lee and K. M. Teo, Multidimensional periodic multiwavelets, *J. Approx. Theory* **98** (1999), 72–103.
- [9] S. S. Goh and V. B. Yap, Matrix extension and biorthogonal multiwavelet construction, *Linear Algebra and Appl.* **269** (1998), 139–157.
- [10] T. N. T. Goodman, Construction of wavelets with multiplicity, *Rendiconti di Matematica* **15** (1994), 665–691.
- [11] Y. W. Koh, S. L. Lee and H. H. Tan, Periodic orthogonal splines and wavelets, *Appl. Comput. Harmonic Anal.* **2** (1995), 201–218.
- [12] F. J. Narcowich and J. D. Ward, Wavelets associated with periodic basis functions, *Appl. Comput. Harmonic Anal.* **3** (1996), 40–56.
- [13] G. Plonka and M. Tasche, On the computation of periodic spline wavelets, *Appl. Comput. Harmonic Anal.* **2** (1995), 1–14.
- [14] I. J. Young, *An Introduction to Nonharmonic Fourier Series* (Academic Press, 1980).

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 KENT RIDGE CRESCENT, SINGAPORE 119260, REPUBLIC OF SINGAPORE

HWA CHONG JUNIOR COLLEGE, 661 BUKIT TIMAH ROAD, SINGAPORE 269734, REPUBLIC OF SINGAPORE