

Solutions to Tutorial 7

Question 1 (a). The limiting function $F(x) = \lim_{n \rightarrow \infty} \frac{n^2}{n^2 + x^2} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{x^2}{n^2}} = 1$.

$$0 \leq T_n = \sup_{0 \leq x \leq 1} |F_n(x) - F(x)| = \sup_{0 \leq x \leq 1} \left| \frac{n^2}{n^2 + x^2} - 1 \right| = \sup_{0 \leq x \leq 1} \frac{x^2}{n^2 + x^2} \leq \frac{1}{n^2}.$$

Since $\lim_{n \rightarrow \infty} \frac{1}{n^2} = \lim_{n \rightarrow \infty} 0 = 0$, we have $\lim_{n \rightarrow \infty} T_n = 0$ by the Squeeze theorem and so $\{F_n\}$ converges uniformly on $[0, 1]$. \square

Question 1 (b). The limiting function $F(x) = \lim_{n \rightarrow \infty} x^n(1 - x) = 0$ for $0 \leq x \leq 1$.

Observe

$$T_n = \sup_{0 \leq x \leq 1} |F_n(x) - F(x)| = \sup_{0 \leq x \leq 1} x^n(1 - x).$$

Let $g(x) = x^n(1 - x)$. Then $g'(x) = nx^{n-1}(1 - x) - x^n = x^{n-1}[n - (n + 1)x]$. From $g'(x) = 0$, we have $x = 0$ or $\frac{n}{n+1}$. Since $g'(x) \geq 0$ for $0 \leq x \leq \frac{n}{n+1}$ and $g'(x) \leq 0$ for $\frac{n}{n+1} \leq x \leq 1$, $\sup_{0 \leq x \leq 1} g(x) = \max\{g(x) | 0 \leq x \leq 1\} = \left(\frac{n}{n+1}\right)^n \left(1 - \frac{n}{n+1}\right)$ and so

$$\lim_{n \rightarrow \infty} T_n = \lim_{n \rightarrow \infty} \left(\frac{n}{n+1}\right)^n \left(1 - \frac{n}{n+1}\right) = \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^n} \cdot \frac{1}{n+1} = \frac{1}{e} \cdot 0 = 0.$$

Thus $\{F_n\}$ converges uniformly on $[0, 1]$. \square

Question 1 (c). The limiting function $f(x) = \lim_{n \rightarrow \infty} \frac{n \ln x}{x^n} = 0$ for $1 \leq x < \infty$. Observe that

$$T_n = \sup_{x \geq 1} |f_n(x) - f(x)| = \sup_{x \geq 1} \left| \frac{n \ln x}{x^n} - 0 \right| = \sup_{x \geq 1} \frac{n \ln x}{x^n}.$$

Let $g(x) = \frac{n \ln x}{x^n}$. From

$$g'(x) = (n \ln x \cdot x^{-n})' = n \frac{1}{x} x^{-n} - n^2 \ln x \cdot x^{-n-1} = \frac{n - n^2 \ln x}{x^{n+1}} = 0,$$

we have $x = e^{\frac{1}{n}}$. Since $g'(x) \geq 0$ for $1 \leq x \leq e^{\frac{1}{n}}$ and $g'(x) \leq 0$ for $x \geq e^{\frac{1}{n}}$, we have

$$T_n = \sup_{x \geq 1} g(x) = \max\{g(x) | x \geq 1\} = \frac{n \ln e^{\frac{1}{n}}}{\left(e^{\frac{1}{n}}\right)^n} = \frac{1}{e}$$

and so $\lim_{n \rightarrow \infty} T_n = \frac{1}{e} \neq 0$. Thus $\{f_n\}$ does not converge uniformly on $[1, +\infty)$. \square

Question 1 (d). The limiting function $f(x) = \lim_{n \rightarrow \infty} \frac{n \ln x \cos nx}{x^n} = 0$ for $x \geq 4$. Observe that

$$0 \leq T_n = \sup_{x \geq 4} |f_n(x) - f(x)| = \sup_{x \geq 4} \frac{n \ln x \cdot |\cos nx|}{x^n} \leq \sup_{x \geq 4} \frac{n \ln x}{x^n} = g(4) = \frac{n \ln 4}{4^n},$$

where $g(x) = \frac{n \ln x}{x^n}$ is monotone decreasing on $[4, +\infty)$. Since $\lim_{n \rightarrow \infty} \frac{n \ln 4}{4^n} = \lim_{n \rightarrow \infty} 0 = 0$, we have $\lim_{n \rightarrow \infty} T_n = 0$ by the Squeeze theorem and so $\{f_n\}$ converges uniformly on $[4, +\infty)$. \square

Question 1 (e). The limiting function $F(x) = \lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} \frac{n^2}{n^2 + x^2} = 1$. Observe that

$$T_n = \sup_{x \geq 0} |F_n(x) - F(x)| = \sup_{x \geq 0} \left| \frac{n^2}{n^2 + x^2} - 1 \right| = \sup_{x \geq 0} \frac{x^2}{n^2 + x^2}.$$

Let $g(x) = \frac{x^2}{n^2 + x^2}$. Since

$$g'(x) = \frac{2x(n^2 + x^2) - x^2 \cdot 2x}{(n^2 + x^2)^2} = \frac{2xn^2}{(n^2 + x^2)^2} \geq 0$$

for $x \geq 0$, the function $g(x)$ is monotone increasing and so

$$T_n = \sup_{x \geq 0} g(x) = \lim_{x \rightarrow \infty} g(x) = \lim_{x \rightarrow \infty} \frac{x^2}{n^2 + x^2} = \lim_{x \rightarrow \infty} \frac{2x}{2x} = 1.$$

Thus $\lim_{n \rightarrow \infty} T_n = 1 \neq 0$ and so $\{F_n(x)\}$ does not converge uniformly on $[0, +\infty)$. \square

Question 2 (i). Since

$$\left| \frac{\cos nx}{n^2 + x^2} \right| \leq \frac{1}{n^2}$$

and the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is convergent by the p -series, the series of functions $\sum_{n=1}^{\infty} \frac{\cos nx}{n^2 + x^2}$ converges uniformly on $(-\infty, +\infty)$ by the Weierstrass M -test. \square

Question 2 (ii). Since

$$\left| \frac{1}{1 + n^3 x^2} \right| \leq \frac{1}{1 + 2^2 n^3} \leq \frac{1}{n^3}$$

for $x \geq 2$ and the series $\sum_{n=1}^{\infty} \frac{1}{n^3}$ is convergent by the p -series, the series of functions

$\sum_{n=1}^{\infty} \frac{1}{1 + n^3 x^2}$ converges uniformly on $[2, +\infty)$ by the Weierstrass M -test. \square

Question 2 (iii). Let $f_n(x) = \frac{xe^{-x}}{n^2}$ for $x \in (0, +\infty)$. Then

$$f'_n(x) = \frac{1}{n^2} (e^{-nx} - xne^{-nx}) = \frac{(1 - nx)e^{-nx}}{n^2}.$$

Thus $f_n(x)$ is monotone increasing for $0 \leq x \leq \frac{1}{n}$ and monotone decreasing for $x \geq \frac{1}{n}$. It follows that

$$|f_n(x)| \leq f_n\left(\frac{1}{n}\right) = \frac{\frac{1}{n}e^{-n \cdot \frac{1}{n}}}{n^2} = \frac{e^{-1}}{n^3}$$

for $x \in (0, +\infty)$. Since the series $\sum_{n=1}^{\infty} \frac{e^{-1}}{n^3} = e^{-1} \sum_{n=1}^{\infty} \frac{1}{n^3}$ is convergent by the p -series, the series of functions $\sum_{n=1}^{\infty} \frac{xe^{-nx}}{n^2}$ converges uniformly on $(0, +\infty)$ by the Weierstrass M -test. \square

Question 3. Since the series of functions $\sum_{n=1}^{\infty} g_n(x)$ converges uniformly on I , by the Cauchy Criterion, for any $\epsilon > 0$, there exists N such that

$$\sum_{k=n+1}^m g_k(x) = \left| \sum_{k=n+1}^m g_k(x) \right| < \epsilon$$

for all $x \in I$ and $m > n > N$. Because $g_k(x) \geq |f_k(x)| \geq 0$, for all $x \in I$ and $n > m > N$, we have

$$\left| \sum_{k=n+1}^m f_k(x) \right| \leq \sum_{k=m+1}^n |f_k(x)| \leq \sum_{k=n+1}^m g_k(x) < \epsilon.$$

Thus, by the Cauchy Criterion, the series of functions $\sum_{n=1}^{\infty} f_n(x)$ converges uniformly. \square

Question 4. Answer: NO. Suppose that the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^x}$ converges uniformly on $(0, +\infty)$. By the Cauchy Criterion, for given $\epsilon = \frac{1}{2}$ there exists N such that

$$\left| \sum_{k=n+1}^m (-1)^{k+1} \frac{1}{k^x} \right| < \frac{1}{2}$$

4

for all $n, m > N$ and $x \in (0, +\infty)$. In particular, let $n = N + 1$ and $m = N + 2$,

$$\frac{1}{(N+2)^x} = \left| \sum_{k=N+2}^{N+2} (-1)^{n+1} \frac{1}{n^x} \right| < \frac{1}{2}$$

for all $x \in (0, +\infty)$. This contradicts to that

$$\frac{1}{(N+2)^x} \geq \frac{1}{2}$$

when $0 < x \leq \frac{\ln 2}{\ln(N+2)}$.

□