

Lecture Notes On Advanced Calculus II

Jie Wu

Department of Mathematics
National University of Singapore

Contents

Chapter 1. Sequences of Real Numbers	5
1. Sequences	5
2. Limits of Sequences	7
3. Sequences which tend to ∞	17
4. Techniques For Computing Limits	19
5. The Least Upper Bounds and the Completeness Property of \mathbb{R}	28
6. Monotone Sequences	39
7. Subsequences	44
8. The Limit Superior and Inferior of a Sequence	47
9. Cauchy Sequences	64
Chapter 2. Series of Real Numbers	71
1. Series	71
2. Tests for Positive Series	82
3. Alternating Series	111
4. Absolute and Conditional Convergence	116
5. Rearrangement properties of a series - an informal treatment	121
6. Remarks on the various tests for convergence/divergence of series	122
Chapter 3. Sequences and Series of Functions	125
1. Pointwise Convergence	125
2. Uniform Convergence	137
3. Uniform Convergence of $\{F_n\}$ and Continuity	156
4. Uniform Convergence and Integration	166
5. Uniform Convergence and Differentiation	173
6. Power Series	177
7. Differentiation of Power Series	198
8. Taylor Series	214
Bibliography	235

Sequences of Real Numbers

1. Sequences

A **sequence** is an ordered list of numbers. For example,

$$1, 2, 3, 4, 5, 6$$

The order of the sequence is important. For example,

$$2, 1, 4, 3, 6, 5$$

is different from above sequence. An **infinite** sequence is a list which does not end. For example,

$$1, 1/2, 1/3, 1/4, 1/5, \dots$$

We are going to study infinite sequences. The Formal Definition of a **(infinite) Sequence** is a function whose domain is the set of positive integers. We denote by $\{a_n\}$ the sequence

$$a_1, a_2, a_3, \dots, a_n, \dots$$

EXAMPLE 1.1. *Here are some examples of infinite sequences.*

- (1). $1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots$
- (2). $\frac{1}{3}, \frac{1}{3^2}, \frac{1}{3^3}, \frac{1}{3^4}, \dots$
- (3). $1, -2, 3, -4, 5, \dots$


Can you find a formula for each of the above sequences?

Answer: (1). $a_n = 1/n$. (2). $a_n = 1/3^n$. (3). $(-1)^{n-1}n$.


Historic Remarks:

While sequences of numbers appear on the earliest of artifacts from prehistory, the study of sequences as abstract patterns of numbers was most likely conducted by the Pythagoreans circa 400 B.C.E. The Pythagoreans generated number sequences by geometrical arrangements of pebbles, called figurate numbers. Although it was a part of their belief that "Everything is number", it led to the birth of the branch of mathematics that is today called number theory. Some examples of figurate numbers include:

The **Triangular Numbers** formed by arranging pebbles in triangular patterns, indicated by the following picture. (The formula is $\frac{n(n+1)}{2}$.)



The **Hexagonal numbers** indicated by the following picture. (The formula is $n(2n - 1)$.)



Links to Other Areas of Science: The topics on sequences are used in many areas of science. For example, much attention has been given to the sequences that are the discrete versions of the logistic differential equation. The recursive equation is simple enough, however the terms of the sequence may exhibit extremely complex behavior. This equation is used as an introduction to dynamical systems and chaos theory a hot area of mathematical research. The **logistic difference equation** is given by

$$p_{n+1} = kp_n(1 - p_n),$$

where $0 < p_0 < 1$ and k is a given constant. In ecology, this is a model for population growth such as modelling insect populations, where mating and death occur in a periodic fashion. For values of k between 3.6 and 4 the behavior becomes chaotic.

2. Limits of Sequences

DEFINITION 2.1. The **limit** of $\{a_n\}$ is A , and is written as


$$\lim_{n \rightarrow \infty} a_n = A,$$

if for any $\epsilon > 0$, there is a natural number N such that for every $n > N$, we have


$$|a_n - A| < \epsilon.$$

Roughly speaking, $\lim_{n \rightarrow \infty} a_n = A$ means that a_n becomes arbitrarily close to A for all sufficiently large n .

Graphically, one has



or



Remark. 1. Some sequences do not satisfy the above. We call such sequences **divergent**.

2. Sequences which satisfy the above definition, i.e. A exists and is finite, are called **convergent** sequences.

EXAMPLE 2.2. *Prove the following limits by using $\epsilon - N$ definition*

- 1) $\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$
- 2) $\lim_{n \rightarrow \infty} \sqrt{\frac{n^2}{n^2 + 1}} = 1.$
- 3) $\lim_{n \rightarrow \infty} \left(\frac{3}{4}\right)^n = 0.$

SOLUTION. (1). $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.

Given any $\epsilon > 0$, we want to find N such that

$$\left| \frac{1}{n} - 0 \right| < \epsilon$$

for $n > N$, i.e.,

$$n > \frac{1}{\epsilon}$$

for $n > N$. Choose N to be the smallest integer such that

$$N \geq \frac{1}{\epsilon}.$$

(N is found now!) When $n > N$, then

$$n > N \geq \frac{1}{\epsilon}$$

or

$$\left| \frac{1}{n} - 0 \right| < \epsilon.$$

Thus $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$.

$$(2). \lim_{n \rightarrow \infty} \sqrt{\frac{n^2}{n^2 + 1}} = 1.$$

Given any $\epsilon > 0$, we want to find N such that

$$\left| \sqrt{\frac{n^2}{n^2 + 1}} - 1 \right| < \epsilon$$

for $n > N$. Now

$$\left| \sqrt{\frac{n^2}{n^2 + 1}} - 1 \right| < \epsilon \iff \left| \frac{n}{\sqrt{n^2 + 1}} - 1 \right| < \epsilon$$

$$\iff \left| \frac{n - \sqrt{n^2 + 1}}{\sqrt{n^2 + 1}} \right| < \epsilon$$

$$\iff \left| \frac{n^2 - (n^2 + 1)}{\sqrt{n^2 + 1}(n + \sqrt{n^2 + 1})} \right| < \epsilon$$

$$\iff \frac{1}{\sqrt{n^2 + 1}(n + \sqrt{n^2 + 1})} < \epsilon$$

$$\iff \sqrt{n^2 + 1}(n + \sqrt{n^2 + 1}) > \frac{1}{\epsilon}$$

Observe that

$$\sqrt{n^2 + 1}(n + \sqrt{n^2 + 1}) > n$$

for $n \geq 1$. Choose N to be the smallest integer such that

$$N \geq \frac{1}{\epsilon}.$$

Then, for $n > N$,

$$\sqrt{n^2 + 1}(n + \sqrt{n^2 + 1}) > \sqrt{N^2 + 1}(N + \sqrt{N^2 + 1})$$

$$> N \geq \frac{1}{\epsilon}$$

or

$$\left| \sqrt{\frac{n^2}{n^2 + 1}} - 1 \right| < \epsilon.$$

Thus N is found and hence the result.

$$(3). \lim_{n \rightarrow \infty} \left(\frac{3}{4}\right)^n = 0.$$

Given any $\epsilon > 0$, we want to find N such that

$$\left| \left(\frac{3}{4}\right)^n - 0 \right| < \epsilon$$

for $n > N$. Observe that

$$\left(\frac{3}{4}\right)^n < \epsilon$$

$$\iff n \ln \left(\frac{3}{4}\right) < \ln(\epsilon)$$

$$\iff n > \frac{\ln(\epsilon)}{\ln(3/4)}$$

(**Note.** $\ln(3/4) < 0$!!) Choose N to be the smallest positive integer such that

$$N \geq \frac{\ln(\epsilon)}{\ln(3/4)}.$$

When $n > N$, then

$$n > N \geq \frac{\ln(\epsilon)}{\ln(3/4)} \quad \text{or}$$

$$\left| \left(\frac{3}{4}\right)^n - 0 \right| < \epsilon.$$

The proof is finished. □

Remark. In the above proofs, we have used the following basic property of real numbers:

Archimedean Property: For any given real number x , there exists a natural number N (depending on x) such that $N > x$.

THEOREM 2.3. *If $\{a_n\}$ has a limit, then the limit is unique.*

PROOF. Let A and B be limits of $\{a_n\}$. Suppose that $A \neq B$. Choose $\epsilon = \frac{|A - B|}{2}$.

Then $\epsilon > 0$ because $A \neq B$. By definition, there exists N_1 and N_2 such that

$$|a_n - A| < \epsilon$$

for $n > N_1$ and

$$|a_n - B| < \epsilon$$

for $n > N_2$.

For $n > \max\{N_1, N_2\}$, we have

$$\begin{aligned} |A - B| &= |(A - a_n) + (a_n - B)| \leq |A - a_n| + |a_n - B| \\ &< 2\epsilon = 2 \frac{|A - B|}{2} = |A - B|, \end{aligned}$$

which is a contradiction. Thus $A = B$. \square

THEOREM 2.4 (Squeeze or Sandwich Theorem). *Given three sequences*

$$\{a_n\}, \{b_n\}, \{c_n\}$$

such that

(i) $a_n \leq b_n \leq c_n$ for every n and

(ii) $\lim_{n \rightarrow \infty} a_n = A = \lim_{n \rightarrow \infty} c_n,$

then $\lim_{n \rightarrow \infty} b_n = A.$

Remark. The above theorem is still applicable if the inequality

$$a_n \leq b_n \leq c_n$$

is true **eventually**.

PROOF. For any $\epsilon > 0$, there exists N_1 and N_2 such that

$$|c_n - A| < \epsilon$$

for $n > N_1$ and

$$|a_n - A| < \epsilon$$

for $n > N_2$. Let $N = \max\{N_1, N_2\}$. For $n > N$, we have

$$-\epsilon < c_n - A < \epsilon \quad \text{and} \quad -\epsilon < a_n - A < \epsilon$$

$$A - \epsilon < c_n < A + \epsilon \quad \text{and} \quad A - \epsilon < a_n < A + \epsilon.$$

Thus

$$A - \epsilon < a_n \leq b_n \leq c_n < A + \epsilon \quad \text{or} \quad |b_n - A| < \epsilon.$$

By definition, we have $\lim_{n \rightarrow \infty} b_n = A$ and hence the result. \square

EXAMPLE 2.5. *Find limits*

$$1) \lim_{n \rightarrow \infty} \frac{1 + \sin n}{n}$$

$$2) \left(\frac{3n - 1}{4n + 1} \right)^n.$$

SOLUTION. (1). Since

$$0 \leq \frac{1 + \sin n}{n} \leq \frac{2}{n}$$

and $\lim_{n \rightarrow \infty} \frac{2}{n} = \lim_{n \rightarrow \infty} 0 = 0$, we have

$$\lim_{n \rightarrow \infty} \frac{1 + \sin n}{n} = 0.$$

(2). Since

$$0 \leq \left(\frac{3n - 1}{4n + 1} \right)^n \leq \left(\frac{3}{4} \right)^n$$

and $\lim_{n \rightarrow \infty} \left(\frac{3}{4} \right)^n = \lim_{n \rightarrow \infty} 0 = 0$, we have

$$\lim_{n \rightarrow \infty} \left(\frac{3n - 1}{4n + 1} \right)^n = 0.$$

□

3. Sequences which tend to ∞

DEFINITION 3.1. $\{a_n\}$ tends to $+\infty$ if for each number k , there is an N such that

$$a_n > k \quad \text{for all } n > N.$$

Remark. For such sequences, we write as $a_n \rightarrow +\infty$ as $n \rightarrow \infty$ or

$$\lim_{n \rightarrow \infty} a_n = +\infty.$$

$\{a_n\}$ tends to $-\infty$ if for each number k , there is an N such that

$$a_n < k \quad \text{for all } n > N.$$

In this case, write $\lim_{n \rightarrow \infty} a_n = -\infty$.

EXAMPLE 3.2. The following sequences tend to $+\infty$

- 1) $a_n = \sqrt{\ln n}$.
- 2) $a_n = (3/2)^n$.

The sequences $-\ln n$, $-n^2$ and etc then tend to $-\infty$.

THEOREM 3.3 (Reciprocal Rule). *Consider a sequence $\{a_n\}$.*

(i) *If $a_n > 0$ for all n and $\lim_{n \rightarrow \infty} \frac{1}{a_n} = 0$, then*

$$\lim_{n \rightarrow \infty} a_n = +\infty.$$

(ii) *If $\lim_{n \rightarrow \infty} a_n = \pm\infty$, then $\lim_{n \rightarrow \infty} \frac{1}{a_n} = 0$.*

PROOF. We only prove (i). For each positive integer k , there exists N such that

$$\left| \frac{1}{a_n} - 0 \right| < \frac{1}{k}$$

for $n > N$ because $\lim_{n \rightarrow \infty} \frac{1}{a_n} = 0$. Then, for $n > N$, $a_n > k$ because $a_n > 0$. This finishes the proof. \square

EXAMPLE 3.4. Since $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$, we have $\lim_{n \rightarrow \infty} \sqrt{n} = \infty$. Similarly, since $\lim_{n \rightarrow \infty} \sqrt{n} = +\infty$, we have $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$.

4. Techniques For Computing Limits

THEOREM 4.1. *Let f be a continuous function. Then*

$$\lim_{n \rightarrow \infty} f(a_n) = f\left(\lim_{n \rightarrow \infty} a_n\right).$$

IDEA OF PROOF. By the definition of continuity, when $x \rightarrow x_0$, $f(x) \rightarrow f(x_0)$. Now $\lim_{n \rightarrow \infty} a_n = A$ means that $a_n \rightarrow A$ when $n \rightarrow \infty$. Thus $f(a_n) \rightarrow f(A)$ when $n \rightarrow \infty$, that is,

$$\lim_{n \rightarrow \infty} f(a_n) = f(A) = f\left(\lim_{n \rightarrow \infty} a_n\right).$$

□

EXAMPLE 4.2.

$$\lim_{n \rightarrow \infty} \sin\left(\frac{n\pi}{2n+1}\right) = \lim_{n \rightarrow \infty} \left(\frac{\pi}{2+1/n}\right) = \sin\left(\frac{\pi}{2}\right) = 1.$$

THEOREM 4.3 (L'Hospital's Rule). *Suppose $a_n = f(n)$, $b_n = g(n)$ for differentiable functions f and g . If $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)}$ is of the form $\frac{\infty}{\infty}$ or $\frac{0}{0}$, then*

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{f'(n)}{g'(n)}.$$

History Remark. Although the theorem is named after Marquis de l'Hospital (1661-1704), it should be called Bernoulli's rule. The story is that in 1691, l'Hospital asked Johann Bernoulli (1667-1748) to provide, for a fee, lectures on the new subject of calculus. L'Hospital subsequently incorporated these lectures into the first calculus text, *L'Analyse des infiniment petits (Analysis of infinitely small quantities)*, published in 1696. The initial version of what is now known as l'Hospital's rule first appeared in this text.

EXAMPLE 4.4. *Show that $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$.*

PROOF.

$$\begin{aligned} \lim_{n \rightarrow \infty} \ln \left[\left(1 + \frac{x}{n}\right)^n \right] &= \lim_{n \rightarrow \infty} n \ln \left(1 + \frac{x}{n}\right) \\ &= \lim_{n \rightarrow \infty} \frac{\ln \left(1 + \frac{x}{n}\right)}{1/n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{1+\frac{x}{n}} \cdot \left(-\frac{x}{n^2}\right)}{-\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{x}{1 + \frac{x}{n}} = x. \end{aligned}$$

Thus $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$. □

THEOREM 4.5. *If $\lim_{n \rightarrow \infty} a_n$ and $\lim_{n \rightarrow \infty} b_n$ exist, then*

- (1). $\{a_n + b_n\}$ converges with $\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$;
- (2). $\{a_n - b_n\}$ converges with $\lim_{n \rightarrow \infty} (a_n - b_n) = \lim_{n \rightarrow \infty} a_n - \lim_{n \rightarrow \infty} b_n$;
- (3). $\{ka_n\}$ converges with $\lim_{n \rightarrow \infty} ka_n = k \lim_{n \rightarrow \infty} a_n$, where k is a fixed constant;
- (4). $\{a_n b_n\}$ converges with $\lim_{n \rightarrow \infty} a_n b_n = \lim_{n \rightarrow \infty} a_n \lim_{n \rightarrow \infty} b_n$;
- (5). $\left\{ \frac{a_n}{b_n} \right\}$ converges with $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$, provided $b_n \neq 0$ and $\lim_{n \rightarrow \infty} b_n \neq 0$.

PROOF. See Bartle-Sherbert [1, page 60-61]. □

EXAMPLE 4.6. *Find the limit of*

$$\ln \left(\frac{n^2 + 3n + 2}{2 + 4n + 2n^2} \right) + \cos \left(\frac{1}{\sqrt{n}} \right).$$

SOLUTION.

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left[\ln \left(\frac{n^2 + 3n + 2}{2 + 4n + 2n^2} \right) + \cos \left(\frac{1}{\sqrt{n}} \right) \right] \\ &= \lim_{n \rightarrow \infty} \left[\ln \left(\frac{(n^2 + 3n + 2)/n^2}{(2 + 4n + 2n^2)/n^2} \right) + \cos \left(\frac{1}{\sqrt{n}} \right) \right] \\ &= \lim_{n \rightarrow \infty} \left[\ln \left(\frac{1 + 3/n + 2/n^2}{2/n^2 + 4/n + 2} \right) + \cos \left(\frac{1}{\sqrt{n}} \right) \right] \\ &= \ln \left(\frac{1 + 0 + 0}{0 + 0 + 2} \right) + \cos 0 = \ln \left(\frac{1}{2} \right) + 1 = 1 - \ln 2. \end{aligned}$$

□

THEOREM 4.7 (Some Standard Limits). *Some standard limits are given as follows.*

1. $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$ for any fixed $p > 0$.

2. $\lim_{n \rightarrow \infty} c^n = 0$ for any fixed c where $|c| < 1$.

3. $\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$ for any fixed $c > 0$.

4. $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$.

5. $\lim_{n \rightarrow \infty} \frac{n^p}{c^n} = 0$ for any fixed p and $c > 1$.

6. $\lim_{n \rightarrow \infty} \frac{c^n}{n!} = 0$ for any fixed c .

7. $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$ for any fixed x .

8. $\lim_{n \rightarrow \infty} \frac{(\ln n)^p}{n^k} = 0$ for any fixed $k > 0$.

PROOF. 1. $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$ for any fixed $p > 0$.

$$\lim_{n \rightarrow \infty} \frac{1}{n^p} = \left(\lim_{n \rightarrow \infty} \frac{1}{n} \right)^p = 0^p = 0.$$

2. $\lim_{n \rightarrow \infty} c^n = 0$ for any fixed c where $|c| < 1$.

Case 1: When $c = 0$, the statement is obvious.

Case 2: When $c > 0$, we have

$$\ln \left(\lim_{n \rightarrow \infty} c^n \right) = \lim_{n \rightarrow \infty} \ln c^n = \lim_{n \rightarrow \infty} n \ln c = -\infty.$$

Thus, $\lim_{n \rightarrow \infty} c^n = 0$.

Case 3: When $c < 0$, we have $-|c|^n \leq c^n \leq |c|^n$ for all n . By Case 2, we have $\lim_{n \rightarrow \infty} (-|c|^n) = 0 = \lim_{n \rightarrow \infty} |c|^n$.

Hence by Squeeze theorem, we also have $\lim_{n \rightarrow \infty} c^n = 0$.

3. $\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$ for any fixed $c > 0$.

$$\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = c^{\lim_{n \rightarrow \infty} \frac{1}{n}} = c^0 = 1.$$

$$4. \lim_{n \rightarrow \infty} \sqrt[n]{n} = 1.$$

$$\begin{aligned} \ln \left(\lim_{n \rightarrow \infty} \sqrt[n]{n} \right) &= \lim_{n \rightarrow \infty} \ln \sqrt[n]{n} \\ &= \lim_{n \rightarrow \infty} \frac{\ln n}{n} = 0 \text{ (by L'Hospital's rule).} \end{aligned}$$

$$\text{Thus, } \lim_{n \rightarrow \infty} \sqrt[n]{n} = e^0 = 1.$$

$$5. \lim_{n \rightarrow \infty} \frac{n^p}{c^n} = 0 \text{ for any fixed } p \text{ and } c > 1.$$

Let k be a fixed positive integer such that $p - k < 0$.
Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{n^p}{c^n} &= \lim_{n \rightarrow \infty} \frac{pn^{p-1}}{c^n \ln c} = \lim_{n \rightarrow \infty} \frac{p(p-1)n^{p-2}}{c^n (\ln c)^2} = \dots \\ &= \lim_{n \rightarrow \infty} \frac{p(p-1) \cdots (p-k+1)n^{p-k}}{c^n (\ln c)^k} \\ &= \lim_{n \rightarrow \infty} \frac{p(p-1) \cdots (p-k+1)}{c^n n^{k-p} (\ln c)^k} = 0 \end{aligned}$$

by L'Hospital's rule.

6. $\lim_{n \rightarrow \infty} \frac{c^n}{n!} = 0$ for any fixed c .

Let $a_n = \frac{c^n}{n!} = \frac{c \cdot c \cdot \dots \cdot c}{n(n-1) \cdot \dots \cdot 1}$. Now fix an integer $M > c$. Then for any $n > M$,

$$0 < a_n = \frac{c \cdot c \cdot \dots \cdot c}{n(n-1) \cdot \dots \cdot (M+1)} a_M < \frac{c}{n} a_M.$$

Note that a_M is a fixed number because M is fixed. Since $\lim_{n \rightarrow \infty} 0 = 0 = \lim_{n \rightarrow \infty} \frac{c}{n} a_M$, by the Squeeze theorem, $\lim_{n \rightarrow \infty} a_n = 0$.

7. $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$ for any fixed x . It was proved in Example 4.4.

8. $\lim_{n \rightarrow \infty} \frac{(\ln n)^p}{n^k} = 0$ for any fixed $k > 0$.

Let $m = \ln n$. Then $n = e^m$. By (5),

$$\lim_{n \rightarrow \infty} \frac{(\ln n)^p}{n^k} = \lim_{m \rightarrow \infty} \frac{m^p}{e^{km}} = \lim_{m \rightarrow \infty} \frac{m^p}{(e^k)^m} = 0,$$

where $e^k > 1$ because $k > 0$. □

Strategy: One can find the limits of many sequences from those of the standard sequences.

EXAMPLE 4.8. *Find the limits*

$$1) \lim_{n \rightarrow \infty} \frac{8^n + (\ln n)^{10} + n!}{n^6 - n!}.$$

$$2) \lim_{n \rightarrow \infty} \left(1 - \frac{1}{2n+1}\right)^{3n}.$$

SOLUTION. (1).

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{8^n + (\ln n)^{10} + n!}{n^6 - n!} &= \lim_{n \rightarrow \infty} \frac{8^n/n! + (\ln n)^{10}/n! + 1}{n^6/n! - 1} \\ &= \frac{0 + 0 + 1}{0 - 1} = -1. \end{aligned}$$

(2).

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(1 - \frac{1}{2n+1}\right)^{3n} &= \lim_{n \rightarrow \infty} \left[\left(1 + \frac{-1}{2n+1}\right)^{2n+1} \right]^{\frac{3n}{2n+1}} \\ &= \lim_{n \rightarrow \infty} \left[\left(1 + \frac{-1}{2n+1}\right)^{2n+1} \right]^{\frac{3}{2+1/n}} = (e^{-1})^{\frac{3}{2}} = \frac{1}{e\sqrt{e}} \end{aligned}$$

□

5. The Least Upper Bounds and the Completeness Property of \mathbb{R}

5.1. From Natural Numbers to Real Numbers. Starting with natural numbers $\mathbb{N} = \{1, 2, 3, \dots\}$, we obtain real numbers \mathbb{R} by adding more and more *new* numbers in the following steps:

Step 1. By adding 0 and negative numbers, we have integers $\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \dots\}$.

Step 2. Then we have rational numbers

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p, q \in \mathbb{Z}, q \neq 0 \right\}.$$

Step 3. Then, by adding **irrational** numbers, we have all real numbers.

Below we give some examples of **irrational** numbers. Recall that a natural number $p > 1$ is called **prime** if p is NOT divisible by any natural numbers other than p and 1. For instance, 2, 3, 5, 7, 11, \dots are primes. Every natural number $n > 1$ admits a unique (prime) **factorization**

$$n = p_1 \cdot p_2 \cdots p_k,$$

where each p_i is prime. For instance, $20 = 2 \cdot 2 \cdot 5$ and $66 = 2 \cdot 3 \cdot 11$.

EXAMPLE 5.1. *If n is a natural number, and there is no natural number whose square is n , then \sqrt{n} is NOT a rational number. In particular, $\sqrt{2}$, $\sqrt{3}$, $\sqrt{5}$, $\sqrt{6}$ are irrational numbers.*

PROOF. Suppose that \sqrt{n} is a rational number. We can write \sqrt{n} as $\frac{a}{b}$, where $a, b \in \mathbb{N}$ and $b \neq 0$. Then

$$\sqrt{n} = \frac{a}{b} \quad \iff \quad n = \frac{a^2}{b^2} \quad \iff \quad a^2 = b^2 n.$$

Any prime occurring in the (unique) factorization of a will occur an even number of times in the factorization of a^2 ; similarly for b and b^2 .

By $a^2 = b^2 n$, any prime that occurs in the factorization of n must occur an even number of times, since all primes occurring in the factorization of $b^2 n$ are exactly those occurring in the factorization of a^2 .

Thus n can be written as

$$n = (p_1 \cdot p_2 \cdots p_k)^2,$$

where, of course, the p_i 's need not be distinct.

This, however, is a contradiction to the hypothesis, since n is expressed as the square of a natural number.

□

5.2. Bounded Sets.

DEFINITION 5.2. A set of real numbers S is **bounded above** if there exists a finite real number M such that

$$x \leq M \quad \forall x \in S.$$

M is called an **upper bound** of S .

DEFINITION 5.3. A set of real numbers S is **bounded below** if there exists a finite real number m such that

$$m \leq x \quad \forall x \in S.$$

m is called a **lower bound** of S .

DEFINITION 5.4. A set which is both bounded above and below is called a **bounded set**.

Remark.

1. Upper bounds and lower bounds are not unique.
2. Some sets only have upper bounds but not lower bounds.
3. Some sets have only lower bounds but not upper bounds.
4. A set which is not bounded is called an **unbounded set**.

EXAMPLE 5.5. *Let*

$$S = \{r \mid r \text{ is a rational number with } r < \sqrt{2}\}.$$

Then S is bounded above.

THEOREM 5.6. *Every convergent sequence is bounded.*

PROOF. Let $\{a_n\}$ be a sequence convergent to A . For $\epsilon = 1$, there exists N such that $|a_n - A| < 1$ or $A - 1 < a_n < A + 1$ for $n > N$.

Choose M and m to be the largest and smallest number of the finite numbers

$$a_1, a_2, \dots, a_N, A + 1, A - 1,$$

respectively.

When $n \leq N$, we have $m \leq a_n \leq M$ because M (m) is the largest (smallest) number of the above finite set. When $n > N$, we have

$$m \leq A - 1 < a_n < A + 1 \leq M.$$

Thus, for all n , we have $m \leq a_n \leq M$ and so $\{a_n\}$ is bounded. The proof is finished. \square

COROLLARY 5.7 (Test for divergence). *If $\{a_n\}$ is unbounded, then $\{a_n\}$ diverges.*

Remark.

1. The converse may not be true, i.e., divergent sequence need not be unbounded.
2. The inverse may not be true, i.e., a bounded sequence may not be convergent.

Example. The sequence $\{1, -1, 1, -1, \dots\}$ is bounded but NOT convergent.

5.3. Infimum and Supremum. Recall that any **finite** set of real numbers has a greatest element (maximum) and a least element (minimum).

EXAMPLE 5.8. $\{-2.5, 3.1, -4.4, 4.5, 5\}$

However, this property does not necessarily hold for infinite sets.

EXAMPLE 5.9. $\{1, 2, 3, 4, \dots, \}$.

DEFINITION 5.10. A real number M ($\neq \pm\infty$) is called the **least upper bound** or **supremum** of a set E if

- (S-i) M is an upper bound of E , i.e., $x \leq M$ for every $x \in E$, and
- (S-ii) if $M' < M$, then M' is not an upper bound of E (i.e., there is an $x \in E$ such that $M' < x$).

We write $M = \sup E$.

Remark.

- (i) $\sup E$ is unique whenever it exists.
- (ii) The main difference between $\sup E$ and $\max E$ is that $\sup E$ **need not** be an element of E , whereas $\max E$ **must be** an element of E if it does exist).
- (iii) If E has a maximum, then $\sup E = \max E$.

EXAMPLE 5.11. 1. Let $E = \{r \in \mathbb{Q} \mid 0 \leq r \leq \sqrt{2}\}$. Then $\sup E = \sqrt{2}$ but $\max E$ does not exist because $\sqrt{2}$ is not a rational number, that is, $\sup E \notin E$.

2. Let $E = \{1/2, 2/3, 3/4, 4/5, 5/6, \dots\}$. Then $\sup E = 1$ and $\max E$ does not exist.

3. Let $E = \{1, 1/2, 1/3, 1/4, 1/5, \dots\}$. Then $\max E = 1 = \sup E$.

DEFINITION 5.12. A real number m ($\neq \pm\infty$) is called the **greatest lower bound** or **infimum** of a set E if

- (i) m is a lower bound of E , i.e., $m \leq x$ for every $x \in E$, and
- (ii) if $m' > m$, then m' is not a lower bound of E (i.e., there exists an $x \in E$ such that $x < m'$).

We write $m = \inf E$.

Remark.

- (i) $\inf E$ is unique whenever it exists.
- (ii) The main difference between $\inf E$ and $\min E$ is that $\inf E$ **need not** be an element of E , whereas $\min E$ must be an element of E if it does exist.
- (iii) If E has a minimum, then $\inf E = \min E$.

EXAMPLE 5.13. 1. Let $E = \{1, 1/2, 1/3, 1/4, \dots, \}$. Then $\inf E = 0$ but $\min E$ does not exist.

2. Let $E = \{r \in \mathbb{Q} \mid 0 \leq r \leq \sqrt{2}\}$. Then $\min E = \inf E = 0$.

5.4. The Completeness of \mathbb{R} . A very basic but important property of the set of real numbers is the following:

THEOREM 5.14 (Completeness Axiom of \mathbb{R}). *The following statement hold for subsets of real numbers:*

- (i) *If E is **bounded above**, then $\sup E$ **exists**.*
- (ii) *If E is **bounded below**, then $\inf E$ **exists**.*


Recall that a set E is bounded if and only if it is bounded above and bounded below. Thus the Completeness Axiom leads to

COROLLARY 5.15. *If E is **bounded**, then **both** $\sup E$ and $\inf E$ **exist**.*

To see this, we will need an important property of rational numbers:

THEOREM 5.16 (Density Theorem of \mathbb{Q}). *For any two **real** numbers $x, y \in \mathbb{R}$ satisfying $x < y$, there exist a **rational** number $r \in \mathbb{Q}$ such that*

$$x < r < y.$$



PROOF. Since $y > x$, we have $y - x > 0$ and thus $\frac{1}{y-x} \in \mathbb{R}$ (and $\frac{1}{y-x} > 0$).

Then by the Archimedean Property of \mathbb{R} , there exist a natural number $n \in \mathbb{N}$ such that

$$n > \frac{1}{y-x} \implies ny - nx > 1 \implies ny > 1 + nx,$$

noting that $y - x > 0$. Then the smallest integer m satisfying $m > nx$ must also satisfy $m \leq nx + 1 < ny$. Thus, we have

$$nx < m < ny \implies x < \frac{m}{n} < y,$$

where $r = m/n$ is rational. \square

EXAMPLE 5.17. Let $E = \{x \in \mathbb{Q} : x^2 < 3\}$. Then $\sup E = \sqrt{3}$.

PROOF. For any $x \in E$, we have

$$x^2 < 3 \implies x < \sqrt{3}$$

upon taking positive square roots on both sides. Thus, E is bounded above (by $\sqrt{3}$). Also, it is easy to see that $E \neq \emptyset$ (for example, $0 \in E$). Thus by the Completeness Axiom of \mathbb{R} , $\sup E$ exists in \mathbb{R} . Since $\sqrt{3}$ is an upper bound for E , it follows from that $\sqrt{3} \geq \sup E$.

Suppose $\sqrt{3} \neq \sup E$. Then we must have $\sqrt{3} > \sup E$. But by the Density Theorem for \mathbb{Q} , there exists a rational number $r \in \mathbb{Q}$ such that

$$\sup E < r < \sqrt{3}.$$

Then $r \in \mathbb{Q}$ and $r^2 < 3$. Hence we have $r \in E$ and $r \leq \sup E$, which contradicts to that $\sup E < r$. Hence we must have $\sqrt{3} = \sup E$. \square

Remark. The above example shows that $\sup E$ may be in \mathbb{R} but NOT in \mathbb{Q} , even though $E \subset \mathbb{Q}$. In other words, the Completeness Axiom does not hold for the set of rational numbers \mathbb{Q} .

We will also adopt the following convention:

DEFINITION 5.18. (i) *If $\emptyset \neq E \subset \mathbb{R}$ is not bounded above, then we write $\sup E = +\infty$.*

(i) *If $\emptyset \neq E \subset \mathbb{R}$ is not bounded below, then we write $\inf E = -\infty$.*

6. Monotone Sequences

DEFINITION 6.1. $\{a_n\}$ is called **monotone increasing (decreasing)** if

$$a_n \leq (\geq) a_{n+1}$$

for every n , that is,

$$a_1 \leq a_2 \leq a_3 \leq a_4 \leq \cdots$$

($a_1 \geq a_2 \geq a_3 \geq \cdots$).

EXAMPLE 6.2. (1). The sequence $\{1/n\}$ is monotone decreasing.

(2). The sequence $\{1/2, 2/3, 3/4, 4/5, 5/6, \cdots\}$ is monotone increasing.


PROPOSITION 6.3. *A monotone increasing (decreasing) sequence is bounded below (above).*

PROOF. Let $\{a_n\}$ be a monotone increasing sequence, that is,

$$a_1 \leq a_2 \leq a_3 \leq \cdots .$$

Then a_1 is a lower bound for $\{a_n\}$ and hence the result. □

Basically only two types of monotone increasing can occur, as illustrated below:



THEOREM 6.4 (Monotone Convergence Theorem).
Let $\{a_n\}$ be a sequence.

- (i) *If $\{a_n\}$ is **monotone increasing** and **bounded above**, then $\{a_n\}$ is **convergent** and*

$$\lim_{n \rightarrow \infty} a_n = \sup_n a_n.$$

- (ii) *If $\{a_n\}$ is **monotone decreasing** and **bounded below**, then $\{a_n\}$ is **convergent** and*

$$\lim_{n \rightarrow \infty} a_n = \inf_n a_n.$$

PROOF. (i). Suppose $\{a_n\}$ is monotone increasing and bounded above.

Then by the Completeness Axiom of \mathbb{R} , $\sup_n a_n$ exists (finite).

Now, given $\epsilon > 0$, since $\sup_n a_n - \epsilon < \sup_n a_n$, it follows that $\sup_n a_n - \epsilon$ is not an upper bound of $\{a_n\}$.

In other words, there exists an integer N such that $a_N > \sup_n a_n - \epsilon$. Then for all $n > N$, we have

$$\sup_n a_n - \epsilon < a_N \leq a_n \leq \sup_n a_n < \sup_n a_n + \epsilon \quad (\text{since } n > N).$$

Equivalently, $\left| a_n - \sup_n a_n \right| < \epsilon$ for all $n > N$ and so $\lim_{n \rightarrow \infty} a_n = \sup_n a_n$ (exists).

The proof of (ii) is similar. □

EXAMPLE 6.5. Let $a_n = \frac{n}{n+1}$, that is,
 $\{a_n\} = \{1/2, 2/3, 3/4, \dots\}$.

Then a_n is monotone increasing and bounded above.
Thus

$$\sup_n a_n = \lim_{n \rightarrow \infty} a_n = 1.$$

COROLLARY 6.6. *If $\{a_n\}$ is monotone increasing (decreasing), then either*

- (i) $\{a_n\}$ is convergent or
- (ii) $\lim_{n \rightarrow \infty} a_n = +\infty(-\infty)$.

PROOF. Suppose $\{a_n\}$ is monotone increasing, then either $\{a_n\}$ is bounded above or not bounded above.

Case (a): If $\{a_n\}$ is bounded above, then by the Monotone Convergence Theorem, $\{a_n\}$ converges.

Case (b): If $\{a_n\}$ is not bounded above, then $\{a_n\}$ has no upper bounds. Thus for any given $k > 0$, k is not an upper bound of $\{a_n\}$. In other words, there exists N such that

$$a_N > k.$$

Since $\{a_n\}$ is monotone increasing, it follows that for all $n > N$,

$$a_n \geq a_N > k.$$

Therefore, $\lim_{n \rightarrow \infty} a_n = +\infty$.

The proof for the case when $\{a_n\}$ is monotone decreasing is similar. \square

7. Subsequences

EXAMPLE 7.1. The following are the subsequences of $\{a_n\} = \{1, -1, 1, -1, 1, -1, \dots\}$.

$$\{a_{2n-1}\} = \{1, 1, 1, \dots\}$$
$$\{a_{2n}\} = \{-1, -1, -1, \dots\}.$$

In general, **subsequences** of $\{a_n\}$ are of the form $\{a_{n_k}\}$, $k = 1, 2, 3, \dots$, with

$$n_1 < n_2 < n_3 < \dots .$$

Note. The rule is that we should choose a_{n_1} first and then a_{n_2} with $n_2 > n_1$ and then a_{n_3} with $n_3 > n_2$, so far and so on (up to infinite). Thus n_1 is at least 1, n_2 is at least 2, n_3 is at least 3, \dots .

THEOREM 7.2. *Suppose $\lim_{n \rightarrow \infty} a_n = A$. Then every subsequence of $\{a_n\}$ also converges to A , that is,*

$$\lim_{k \rightarrow \infty} a_{n_k} = A.$$

PROOF. For any given $\epsilon > 0$, since $\lim_{n \rightarrow \infty} a_n = A$, there exists N such that

$$|a_n - A| < \epsilon \quad \text{for all } n > N.$$

Then for all $k > N$, we have

$$n_k \geq k > N.$$

Hence

$$|a_{n_k} - A| < \epsilon \quad \text{for all } k > N.$$

Therefore, $\lim_{k \rightarrow \infty} a_{n_k} = A$. □

COROLLARY 7.3. *Suppose that $\{a_n\}$ has two subsequences that converge to different limits. Then $\{a_n\}$ is divergent.* \square

EXAMPLE 7.4. *The sequence $\{1, -1, 1, -1, \dots\}$ is divergent because $\{a_{2n-1}\} = \{1, 1, \dots\}$ converges to 1 and $\{a_{2n}\} = \{-1, -1, \dots\}$ converges to -1 .*

8. The Limit Superior and Inferior of a Sequence

Given a sequence $\{a_n\}$, we can form another sequence $\{b_n\}$ given by

$$b_n = \sup_{k \geq n} a_k = \sup\{a_n, a_{n+1}, a_{n+2}, \dots\}.$$

EXAMPLE 8.1. Let $\{a_n\} = \{1, -1, 1, -1, \dots\}$. Then

$$b_n = \sup_{k \geq n} a_k = \sup\{\pm 1, \mp 1, \pm 1, \mp 1, \dots\} = 1.$$

PROPOSITION 8.2. For any sequence $\{a_n\}$, the associated sequence $\{b_n\} = \{\sup_{k \geq n} a_k\}$ is always monotone decreasing.

PROOF. For each n ,

$$\begin{aligned} b_n &= \sup\{a_n, a_{n+1}, a_{n+2}, \dots\} \\ &\geq \sup\{a_{n+1}, a_{n+2}, \dots\} = b_{n+1}. \end{aligned}$$

□

DEFINITION 8.3. The **limit superior** of $\{a_n\}$, denoted by $\limsup a_n$ or $\limsup_{n \rightarrow \infty} a_n$ or $\overline{\lim}_{n \rightarrow \infty} a_n$ is defined to be $\lim_{n \rightarrow \infty} b_n$, i.e.

$$\overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \sup_{k \geq n} a_k.$$

EXAMPLE 8.4. 1. Let $\{a_n\} = \{1, -1, 1, -1, 1, -1, \dots\}$.

$$\overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} 1 = 1.$$

2. Let $\{a_n\} = \{1, 2, 3, \dots\}$. Then

$$b_n = \sup_{k \geq n} a_k = \sup\{n, n+1, \dots\} = +\infty$$

and so $\overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = +\infty$.

3. Let $\{a_n\} = \{-1, -2, -3, \dots\}$. Then

$$b_n = \sup_{k \geq n} a_k = \sup\{-n, -n-1, \dots\} = -n$$

and so $\overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = -\infty$.

THEOREM 8.5. *Given any sequence $\{a_n\}$, either*

(1). $\overline{\lim}_{n \rightarrow \infty} a_n$ *exists (finite), or*

(2). $\overline{\lim}_{n \rightarrow \infty} a_n = +\infty$, **or**

(3). $\overline{\lim}_{n \rightarrow \infty} a_n = -\infty$.

PROOF. If $\{a_n\}$ is not bounded above, then each b_n is $+\infty$, and thus

$$\overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = +\infty.$$

If $\{a_n\}$ is bounded above, then each b_n is finite. Since $\{b_n\}$ is monotone decreasing, by Corollary 1.6.6, $\{a_n\}$ converges (to a finite limit), or $\lim_{n \rightarrow \infty} b_n = -\infty$. \square

Similarly, given any sequence $\{a_n\}$, we can form another sequence $\{c_n\}$ given by

$$c_n = \inf_{k \geq n} a_k = \inf \{a_n, a_{n+1}, a_{n+2}, \dots\}.$$

DEFINITION 8.6. The **limit inferior** of $\{a_n\}$, denoted by $\liminf a_n$ or $\liminf_{n \rightarrow \infty} a_n$ or $\underline{\lim}_{n \rightarrow \infty} a_n$ is defined to be $\lim_{n \rightarrow \infty} c_n$, i.e.

$$\underline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \inf_{k \geq n} a_k.$$

EXAMPLE 8.7. 1. Let $\{a_n\} = \{1, -1, 1, -1, 1, -1, \dots\}$.

$$\begin{aligned} \liminf_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} (\inf\{\pm 1, \mp 1, \pm 1, \mp 1, \dots\}) \\ &= \lim_{n \rightarrow \infty} -1 = -1. \end{aligned}$$

2. Let $\{a_n\} = \{1, 2, 3, \dots\}$. Then

$$c_n = \inf_{k \geq n} a_k = \inf\{n, n+1, \dots\} = n$$

and so $\liminf_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = +\infty$.

3. Let $\{a_n\} = \{-1, -2, -3, \dots\}$. Then

$$c_n = \inf_{k \geq n} a_k = \inf\{-n, -n-1, \dots\} = -\infty$$

and so $\liminf_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = -\infty$.

PROPOSITION 8.8. (i). *As in Proposition 8.2, for any given sequence $\{a_n\}$, the associated sequence $\{c_n\} = \{\inf_{k \geq n} a_k\}$ is always monotone increasing.*

(ii). *As in Theorem 8.5, for any given $\{a_n\}$, $\liminf_{n \rightarrow \infty} a_n$ either exists (finite), or $+\infty$, or $-\infty$.*

Remark. We always have

$$\liminf_{n \rightarrow \infty} a_n \leq \overline{\lim}_{n \rightarrow \infty} a_n$$

because $c_n \leq b_n$.

PROPOSITION 8.9. (i). If $\overline{\lim}_{n \rightarrow \infty} a_n = B$ with $B \neq -\infty$, then given $\epsilon > 0$, there exists N such that

$$a_n < B + \epsilon$$

for all $n > N$.

(ii). $\underline{\lim}_{n \rightarrow \infty} a_n = C$ with $C \neq +\infty$, then given $\epsilon > 0$, there exists N such that

$$a_n > C - \epsilon$$

for all $n > N$.

PROOF. (i). If $B = +\infty$, the assertion is obvious and so we assume that B is finite.

Since $\overline{\lim}_{n \rightarrow \infty} a_n = B$, given any $\epsilon > 0$, there exists N such that for all $n > N$,


$$|b_n - B| < \epsilon \implies b_n < B + \epsilon$$

$$\implies \sup\{a_n, a_{n+1}, \dots\} < B + \epsilon,$$


i.e. $a_n, a_{n+1}, \dots < B + \epsilon$ for all $n > N$.

Proof of (ii) is similar. □

Remark. Roughly speaking, Proposition 8.9 says that for any sequence $\{a_n\}$, the a_n 's are eventually $\geq \underline{\lim}_{n \rightarrow \infty} a_n$ and $\leq \overline{\lim}_{n \rightarrow \infty} a_n$ (up to an arbitrarily small quantity ϵ).



In terms of subsequences of $\{a_n\}$, $\overline{\lim}_{n \rightarrow \infty} a_n$ is the **largest (subsequential) limit** of the convergent subsequences of $\{a_n\}$, including those possible subsequences tending to $+\infty$ or $-\infty$. Similarly, $\underline{\lim}_{n \rightarrow \infty} a_n$ is the **smallest subsequential limit**.



This is described in the following theorem.

THEOREM 8.10. *Let $\{a_n\}$ be any sequence. Let $B = \overline{\lim}_{n \rightarrow \infty} a_n$ and let $C = \underline{\lim}_{n \rightarrow \infty} a_n$.*

(i) *Let $\{a_{n_k}\}$ be any subsequence of $\{a_n\}$ such that $\lim_{k \rightarrow \infty} a_{n_k}$ exists, $+\infty$, or $-\infty$. Then*

$$C = \underline{\lim}_{n \rightarrow \infty} a_n \leq \lim_{k \rightarrow \infty} a_{n_k} \leq \overline{\lim}_{n \rightarrow \infty} a_n = B.$$

(ii) **There exists a subsequence $\{a_{n_k}\}$ of $\{a_n\}$ such that**

$$\lim_{k \rightarrow \infty} a_{n_k} = B.$$

(iii) **There exists a subsequence $\{a_{m_k}\}$ of $\{a_n\}$ such that**

$$\lim_{k \rightarrow \infty} a_{m_k} = C.$$

PROOF. The proof is omitted. See the packed lecture notes. □

EXAMPLE 8.11. *Find the limit inferior and limit superior of the following sequences*

- i) $\left\{ \frac{1 - 2(-1)^n n}{3n + 2} \right\},$
- ii) $\left\{ (1 + (-1)^n) \sin \frac{n\pi}{4} \right\},$
- iii) $\{[1.5 + (-1)^n]^n\}.$

SOLUTION. (i). Note that

$$a_{2k} = \frac{1 - 2 \cdot 2k}{3 \cdot 2k + 2}$$

$$\lim_{k \rightarrow \infty} a_{2k} = \lim_{k \rightarrow \infty} \frac{1/k - 4}{6 + 2/k} = -\frac{4}{6} = -\frac{2}{3}$$

$$a_{2k-1} = \frac{1 + 2 \cdot (2k - 1)}{3 \cdot (2k - 1) + 2}$$

$$\lim_{k \rightarrow \infty} a_{2k-1} = \lim_{k \rightarrow \infty} \frac{1/k + 2 \cdot (2 - 1/k)}{3 \cdot (2 - 1/k) + 2/k} = \frac{4}{6} = \frac{2}{3}$$

Thus the subsequential limits are $\pm \frac{2}{3}$ and so

$$\overline{\lim}_{n \rightarrow \infty} a_n = \frac{2}{3} \quad \text{and} \quad \underline{\lim}_{n \rightarrow \infty} a_n = -\frac{2}{3}.$$

(ii). The sequence

$$\left\{ (1 + (-1)^n) \sin \frac{n\pi}{4} \right\}$$

$$= \left\{ 0, 2 \sin \frac{2\pi}{4} = 2, 0, 2 \sin \frac{4\pi}{4} = 0, 0, 2 \sin \frac{6\pi}{4} = -2, \right.$$

$$\left. 0, 2 \sin \frac{8\pi}{4} = 0, 0, 2 \sin \frac{10\pi}{4} = 2, \dots \right\}.$$

The subsequential limits are -2 , 0 and 2 . Thus

$$\overline{\lim}_{n \rightarrow \infty} a_n = 2 \quad \text{and} \quad \underline{\lim}_{n \rightarrow \infty} a_n = -2.$$

(iii). $\{[1.5 + (-1)^n]^n\}$.

Note that

$$a_{2k} = 2.5^{2k} \quad \lim_{k \rightarrow \infty} a_{2k} = \lim_{k \rightarrow \infty} (2.5^k)^2 = +\infty$$

$$a_{2k-1} = 0.5^{2k-1}$$

$$\lim_{k \rightarrow \infty} a_{2k-1} = \lim_{k \rightarrow \infty} \left[\left(\frac{1}{2} \right)^k \right]^{(2k-1)/k}$$

$$= \lim_{k \rightarrow \infty} \left[\left(\frac{1}{2} \right)^k \right]^{2-1/k} = 0^2 = 0.$$

The subsequential limits are 0 and $+\infty$. Thus

$$\overline{\lim}_{n \rightarrow \infty} a_n = +\infty \quad \text{and} \quad \underline{\lim}_{n \rightarrow \infty} a_n = 0.$$

□

The following important theorem was originally proved by Bernhard Bolzano (1781-1848) and modified slightly by Karl Weierstrass (1815-1897). The Bolzano-Weierstrass Theorem will be mentioned again in applied mathematics courses such as MA 3236, Nonlinear Programming.

COROLLARY 8.12 (Bolzano-Weierstrass Theorem).
Every bounded sequence has a convergent subsequence.

PROOF. Let $\{a_n\}$ be a bounded sequence. Since $\{a_n\}$ is bounded,

$$\begin{aligned} -\infty < \inf\{a_1, a_2, \dots\} = c_1 &\leq \underline{\lim}_{n \rightarrow \infty} a_n \\ &\leq \overline{\lim}_{n \rightarrow \infty} a_n \leq b_1 = \sup\{a_1, a_2, \dots\} < +\infty. \end{aligned}$$

Thus $\overline{\lim}_{n \rightarrow \infty} a_n$ is finite.

By Part (ii) of Theorem 8.10, there exists a convergent subsequence $\{a_{n_k}\}$ of $\{a_n\}$ such that

$$\lim_{k \rightarrow \infty} a_{n_k} = \overline{\lim}_{n \rightarrow \infty} a_n.$$

□

THEOREM 8.13. $\liminf_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n$ (*finite, $+\infty$, $-\infty$*) **if and only if** $\lim_{n \rightarrow \infty} a_n$ *exists (finite), $+\infty$, or $-\infty$.*

PROOF. Suppose that $\liminf_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n = A$. Let $b_n = \sup\{a_n, a_{n+1}, \dots\}$ and let $c_n = \inf\{a_n, a_{n+1}, \dots\}$.

Then

$$c_n = \inf\{a_n, a_{n+1}, \dots\} \leq a_n \leq b_n = \sup\{a_n, a_{n+1}, \dots\}.$$

By the assumption, we have

$$\lim_{n \rightarrow \infty} c_n = \liminf_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n.$$

By the Squeeze theorem, the sequence $\{a_n\}$ converges and

$$\lim_{n \rightarrow \infty} a_n = \liminf_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n.$$

Conversely suppose that $\{a_n\}$ converges, tends to $+\infty$, or tends to $-\infty$. Let $A = \lim_{n \rightarrow \infty} a_n$ and let $\{a_{n_k}\}$ be any subsequence of $\{a_n\}$.

By Theorem 7.2, we have $\lim_{k \rightarrow \infty} a_{n_k} = A$. Thus the only subsequential limit of $\{a_n\}$ is A .

By Theorem 8.10, we have

$$\liminf_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n = A = \lim_{n \rightarrow \infty} a_n.$$

□

Remark. This theorem means that

1) If $\underline{\lim}_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n$, then

$$\lim_{n \rightarrow \infty} a_n = \underline{\lim}_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n.$$

2) If $\lim_{n \rightarrow \infty} a_n$ exists, $+\infty$ or $-\infty$, then

$$\underline{\lim}_{n \rightarrow \infty} a_n = \overline{\lim}_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} a_n.$$

PROPOSITION 8.14. *Let $a_n > 0$ for all n . Prove that*

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq \overline{\lim}_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}.$$

PROOF. Let $B = \overline{\lim}_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$. If $B = +\infty$, clearly

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq B = +\infty.$$

So we may assume that $B < +\infty$.

Since $a_n > 0$, we have $\frac{a_{n+1}}{a_n} > 0$ and so $B \geq 0$. Thus B is a finite nonnegative number.

By Proposition 8.9, given any $\epsilon > 0$, there exists N such that

$$\frac{a_{n+1}}{a_n} < B + \epsilon$$

for $n > N$.

Fixed any n with $n > N$, we have

$$0 < \frac{a_{n+1}}{a_n}, \frac{a_{n+2}}{a_{n+1}}, \frac{a_{n+3}}{a_{n+2}}, \dots < B + \epsilon$$

and so, for any $k \geq 1$, we have

$$\begin{aligned} 0 < \frac{a_{n+1}}{a_n} \cdot \frac{a_{n+2}}{a_{n+1}} \cdot \frac{a_{n+3}}{a_{n+2}} \cdots \frac{a_{n+k}}{a_{n+k-1}} &= \frac{a_{n+k}}{a_n} \leq (B + \epsilon)^k \\ &\implies a_{n+k} \leq a_n (B + \epsilon)^k \\ \implies a_{n+k}^{\frac{1}{n+k}} &\leq a_n^{\frac{1}{n+k}} \cdot (B + \epsilon)^{\frac{k}{n+k}} \quad \text{for any } k \geq 1. \end{aligned}$$

Let k tends to ∞ . We have

$$\begin{aligned} & \lim_{k \rightarrow \infty} a_n^{\frac{1}{n+k}} \cdot (B + \epsilon)^{\frac{k}{n+k}} \\ = & \lim_{k \rightarrow \infty} a_n^{\frac{1}{n+k}} \cdot \lim_{k \rightarrow \infty} (B + \epsilon)^{\frac{1}{n/k+1}} = a_n^0 \cdot (B + \epsilon) = B + \epsilon. \end{aligned}$$

Thus

$$\begin{aligned} \overline{\lim}_{m \rightarrow \infty} a_m^{\frac{1}{m}} &= \overline{\lim}_{k \rightarrow \infty} a_{n+k}^{\frac{1}{n+k}} \leq \overline{\lim}_{k \rightarrow \infty} a_n^{\frac{1}{n+k}} \cdot (B + \epsilon)^{\frac{k}{n+k}} \\ &= \lim_{k \rightarrow \infty} a_n^{\frac{1}{n+k}} \cdot (B + \epsilon)^{\frac{k}{n+k}} = B + \epsilon. \end{aligned}$$

In other words,

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq B + \epsilon$$

for any $\epsilon > 0$

and so

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{\epsilon \rightarrow 0} \left(\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \right) \leq \lim_{\epsilon \rightarrow 0} (B + \epsilon) = B,$$

that is,

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}.$$

□

EXERCISE 8.1. *Let $a_n > 0$ for all n . Prove that*

$$\underline{\lim}_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} \leq \underline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

By Proposition 8.14 and Exercise 8.1, we have

$$\underline{\lim}_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} \leq \underline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} \leq \overline{\lim}_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}.$$

PROPOSITION 8.15. *Let $a_n > 0$ for all n . **Suppose that the limit $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$ exists or $+\infty$. Then $\lim_{n \rightarrow \infty} \sqrt[n]{a_n}$ exists or $+\infty$, with***

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}.$$

For instance,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\sqrt[n]{n!}}{n} &= \lim_{n \rightarrow \infty} \sqrt[n]{\frac{n!}{n^n}} = \lim_{n \rightarrow \infty} \frac{(n+1)!}{(n+1)^{n+1}} \cdot \frac{n^n}{n!} \\ &= \lim_{n \rightarrow \infty} \frac{1}{(n+1)^n/n^n} = \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^n} = \frac{1}{e}. \end{aligned}$$


9. Cauchy Sequences

DEFINITION 9.1. $\{a_n\}$ is called a **Cauchy sequence** if given any $\epsilon > 0$, there exists a natural number N such that for all $m, n > N$, we have

$$|a_n - a_m| < \epsilon.$$

Remark.

Roughly speaking, a sequence is Cauchy if the width of its tail $\rightarrow 0$ as $n \rightarrow \infty$.



PROPOSITION 9.2. Every Cauchy sequence is bounded.

PROOF. Let $\{a_n\}$ be a Cauchy sequence. Choose $\epsilon = 1$. There exists N such that $|a_n - a_m| < 1$ for $n, m > N$. In particular, $|a_n - a_{N+1}| < 1$ or

$$a_{N+1} - 1 < a_n < a_{N+1} + 1$$

for $n > N$.

Let

$$M = \max\{a_1, a_2, \dots, a_N, a_{N+1} + 1\}$$
$$m = \min\{a_1, a_2, \dots, a_N, a_{N+1} - 1\}.$$

For $n \leq N$, we have $m \leq a_n \leq M$, and, for $n > N$, we have

$$m \leq a_{N+1} - 1 < a_n < a_{N+1} + 1 \leq M.$$

Thus, for all n , we have $m \leq a_n \leq M$ and so $\{a_n\}$ is bounded. \square

The following Criterion was formulated by Augustin-Louis Cauchy (1789-1857).

THEOREM 9.3 (Cauchy's criterion). **A sequence is a convergent sequence if and only if it is a Cauchy sequence.**

PROOF. \implies , i.e., every convergent sequence is Cauchy.

Given that $\{a_n\}$ is convergent, say $\lim_{n \rightarrow \infty} a_n = A$. Then for any given $\epsilon > 0$, there exists N such that

$$|a_n - A| < \frac{\epsilon}{2}$$

for all $n > N$. Now for any $m, n > N$,

$$\begin{aligned} |a_n - a_m| &= |(a_n - A) - (a_m - A)| \leq |a_n - A| + |a_m - A| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

since both $m, n > N$. Therefore, $\{a_n\}$ is a Cauchy sequence.

\Leftarrow , i.e., every Cauchy sequence is convergent.

Given that $\{a_n\}$ is Cauchy. By Proposition 9.2, $\{a_n\}$ is bounded.

By the Bolzano-Weierstrass Theorem (Corollary 8.12), there exists a convergent subsequence $\{a_{n_k}\}$ of $\{a_n\}$. Let $A = \lim_{k \rightarrow \infty} a_{n_k}$.

Given any $\epsilon > 0$, since $\{a_n\}$ is Cauchy, there exists N_1 such that

$$|a_n - a_m| < \frac{\epsilon}{2} \quad \text{for all } n, m > N_1.$$

Since $\{a_{n_k}\}$ converges to A , there exists K such that

$$|a_{n_k} - A| < \frac{\epsilon}{2} \quad \text{for all } k > K.$$

Let $N = \max\{K, N_1\}$. Choose an n_k such that $k > N$, for instance, choose n_k to be n_{N+1} . When $n > N$, by triangular inequality,

$$\begin{aligned} |a_n - A| &= |(a_n - a_{n_k}) + (a_{n_k} - A)| \leq |a_n - a_{n_k}| + |a_{n_k} - A| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

because $n > N \geq N_1$, $n_k \geq k > N \geq N_1$ and $k > N \geq K$.

Therefore $\{a_n\}$ converges to A by the definition. \square

EXAMPLE 9.4. Let $s_n = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{n^2}$.
Show that $\{s_n\}$ is convergent.

PROOF. For each $k \geq 1$, we have

$$\begin{aligned}
 & |s_{n+k} - s_n| \\
 = & \left| \left(1 + \frac{1}{2^2} + \cdots + \frac{1}{n^2} + \frac{1}{(n+1)^2} + \cdots + \frac{1}{(n+k)^2} \right) \right. \\
 & \left. - \left(1 + \frac{1}{2^2} + \cdots + \frac{1}{n^2} \right) \right| \\
 = & \frac{1}{(n+1)^2} + \frac{1}{(n+2)^2} + \cdots + \frac{1}{(n+k)^2} \\
 \leq & \frac{1}{n(n+1)} + \frac{1}{(n+1)(n+2)} + \frac{1}{(n+2)(n+3)} \\
 & + \cdots + \frac{1}{(n+k-1)(n+k)} \\
 = & \left(\frac{1}{n} - \frac{1}{n+1} \right) + \left(\frac{1}{n+1} - \frac{1}{n+2} \right) \\
 & + \cdots + \left(\frac{1}{n+k-1} - \frac{1}{n+k} \right) \\
 = & \frac{1}{n} - \frac{1}{n+k} < \frac{1}{n}.
 \end{aligned}$$

Given any $\epsilon > 0$, choose N such that $\frac{1}{N} < \epsilon$, that is, $N > \frac{1}{\epsilon}$. When $m > n > N$, from the above,

$$|s_m - s_n| < \frac{1}{n} < \frac{1}{N} < \epsilon.$$

Thus $\{s_n\}$ is a Cauchy sequence and hence $\{s_n\}$ converges by the Cauchy Criterion. \square

Series of Real Numbers

1. Series

The notion of series is closely related to the sum of numbers. In fact, whenever one hears the word series, the first thing to come to mind is the sum of numbers. This is the basic difference between series and sequences.

The expression

$$a_1 + a_2 + a_3 + \cdots$$

written alternatively as $\sum_{k=1}^{\infty} a_k$ is called an **infinite series**.

EXAMPLE 1.1. (1). $1 + 2 + 3 + 4 + \cdots$.

(2). $1 + 1/2 + 1/3 + 1/4 + \cdots$

(3). $1 + 1/2^2 + 1/3^2 + 1/4^2 + \cdots$.

(4). $1 + 0 + 1 + 0 + 1 + 0 + \cdots$.

DEFINITION 1.2. Given a series $\sum_{k=1}^{\infty} a_k$, its n^{th} **partial sum** S_n is given by

$$S_n = \sum_{k=1}^n a_k = a_1 + a_2 + \cdots + a_n.$$

The sequence $\{S_n\}$ is called the **sequence of partial sums** of the series $\sum_{k=1}^{\infty} a_k$.

For instance,

$$\begin{aligned} S_1 &= a_1 \\ S_2 &= a_1 + a_2 \\ S_3 &= a_1 + a_2 + a_3 \\ S_4 &= a_1 + a_2 + a_3 + a_4 \\ &\dots \end{aligned}$$

EXAMPLE 1.3. Consider the series $1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \cdots$. The $S_{2n-1} = 1$ and $S_{2n} = 0$.

DEFINITION 1.4. Consider the sequence of partial sums $\{S_n\}$ of the series $\sum_{k=1}^{\infty} a_k$. If this sequence converges to a number S , we say that the series $\sum_{k=1}^{\infty} a_k$ **converges** to S and write

$$\sum_{k=1}^{\infty} a_k = \lim_{n \rightarrow \infty} S_n = S.$$

If $\{S_n\}$ diverges, then we say $\sum_{k=1}^{\infty} a_n$ **diverges**.

EXAMPLE 1.5 (Geometric Series). Let $a \neq 0$. Consider the series

$$\sum_{n=0}^{\infty} ar^n = a + ar + ar^2 + ar^3 + \cdots .$$

Then the partial sum

$$\begin{aligned} S_n &= a + ar + ar^2 + \cdots + ar^{n-1} = a(1 + r + \cdots + r^{n-1}) \\ &= \begin{cases} a \frac{1 - r^n}{1 - r} & r \neq 1 \\ an & r = 1 \end{cases} \end{aligned}$$

When $-1 < r < 1$, $S_n \rightarrow \frac{a}{1 - r}$ as $n \rightarrow \infty$.

When $r > 1$, S_n diverges because $r^n \rightarrow +\infty$ as $n \rightarrow \infty$.

When $r = 1$, $S_n = an$ diverges.

When $r = -1$, $S_n = \frac{a[1 - (-1)^n]}{2}$ diverges.

When $r < -1$, S_n diverges because $r^n \rightarrow \pm\infty$.

Thus **the geometric series** $\sum_{n=0}^{\infty} ar^n$ **converges if and only if** $-1 < r < 1$, **and,**

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1 - r}$$

for $-1 < r < 1$.

Remark. If $\sum_{k=1}^{\infty} a_k$ and $\sum_{k=1}^{\infty} b_k$ converges, then one always has

$$(i) \quad \sum_{k=1}^{\infty} (a_k + b_k) = \sum_{k=1}^{\infty} a_k + \sum_{k=1}^{\infty} b_k.$$

$$(ii) \quad \sum_{k=1}^{\infty} ca_k = c \sum_{k=1}^{\infty} a_k.$$

EXAMPLE 1.6.

$$\begin{aligned} \sum_{n=1}^{\infty} \left[\left(\frac{1}{4}\right)^n + \left(\frac{1}{5}\right)^n \right] &= \sum_{n=1}^{\infty} \left(\frac{1}{4}\right)^n + \sum_{n=1}^{\infty} \left(\frac{1}{5}\right)^n \\ &= \left[\frac{1}{4} + \left(\frac{1}{4}\right)^2 + \dots \right] + \left[\frac{1}{5} + \left(\frac{1}{5}\right)^2 + \dots \right] \\ &= \frac{1}{4} \left[1 + \frac{1}{4} + \left(\frac{1}{4}\right)^2 + \dots \right] + \frac{1}{5} \left[1 + \frac{1}{5} + \left(\frac{1}{5}\right)^2 + \dots \right] \\ &= \frac{1}{4} \frac{1}{1 - \frac{1}{4}} + \frac{1}{5} \frac{1}{1 - \frac{1}{5}} \\ &= \frac{1}{4 \cdot \frac{3}{4}} + \frac{1}{5 \cdot \frac{4}{5}} = \frac{1}{3} + \frac{1}{4} = \frac{7}{12}. \end{aligned}$$

THEOREM 1.7. If $\sum_{k=1}^{\infty} a_k$ converges, then $\lim_{k \rightarrow \infty} a_k = 0$.

PROOF. Recall that partial sum $S_k = a_1 + a_2 + \cdots + a_k$. We have

$$\begin{aligned} S_k - S_{k-1} &= (a_1 + a_2 + \cdots + a_{k-1} + a_k) \\ &\quad - (a_1 + a_2 + \cdots + a_{k-1}) = a_k. \end{aligned}$$

Since the series $\sum_{k=1}^{\infty} a_k$ converges, the sequence $\{S_k\}$ converges. Let $S = \lim_{k \rightarrow \infty} S_k$. Then

$$\begin{aligned} \lim_{k \rightarrow \infty} a_k &= \lim_{k \rightarrow \infty} (S_k - S_{k-1}) \\ &= \lim_{k \rightarrow \infty} S_k - \lim_{k \rightarrow \infty} S_{k-1} = S - S = 0. \end{aligned}$$

□

COROLLARY 1.8 (Divergence Test). If $\lim_{n \rightarrow \infty} a_n \neq 0$ (or does not exist), then $\sum_{n=1}^{\infty} a_n$ diverges. □

EXAMPLE 1.9. (1). The series $\sum_{n=1}^{\infty} (-1)^n$ is divergent because the limit of the n -th term $(-1)^n$ does not exist.

(2). The series $\sum_{n=1}^{\infty} \frac{n!}{n^2}$ is divergent because

$$\lim_{n \rightarrow \infty} \frac{n!}{n^2} = \frac{1}{\lim_{n \rightarrow \infty} \frac{n^2}{n!}} = \frac{1}{0} = +\infty \neq 0.$$

(3). The series $\sum_{n=1}^{\infty} \frac{2n+1}{3n+2}$ is divergent because

$$\lim_{n \rightarrow \infty} \frac{2n+1}{3n+2} = \lim_{n \rightarrow \infty} \frac{2+1/n}{3+2/n} = \frac{2}{3} \neq 0.$$

Remark. The divergence test is a “one-way” test, i.e., $\lim_{n \rightarrow \infty} a_n = 0$ does NOT imply $\sum_{n=1}^{\infty} a_n$ converges.

THEOREM 1.10 (Cauchy Criterion). *The series $\sum_{k=1}^{\infty} a_k$ converges if and only if given any $\epsilon > 0$, there exists N such that*

$$\left| \sum_{k=n+1}^m a_k \right| < \epsilon$$

for all $m > n > N$.

PROOF. The series $\sum_{k=1}^{\infty} a_k$ converges if and only if the sequence of its partial sums $\{S_n\}$ converges, (by definition), if and only if $\{S_n\}$ is Cauchy. The result follows from

$$\begin{aligned} |S_m - S_n| &= |(a_1 + a_2 + \cdots + a_n + a_{n+1} + \cdots + a_m) \\ &\quad - (a_1 + a_2 + \cdots + a_n)| \\ &= |a_{n+1} + \cdots + a_m| = \left| \sum_{k=n+1}^m a_k \right|. \end{aligned}$$

□

EXAMPLE 1.11 (Harmonic Series). *Show that the series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.*

PROOF. Note that

$$\begin{aligned} \left| \sum_{k=n+1}^{2n} \frac{1}{k} \right| &= \left| \frac{1}{n+1} + \cdots + \frac{1}{2n} \right| \\ &= \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \\ &\geq \frac{1}{2n} + \frac{1}{2n} + \cdots + \frac{1}{2n} = n \cdot \frac{1}{2n} = \frac{1}{2}. \end{aligned}$$

Suppose that the series $\sum_{n=1}^{\infty} \frac{1}{n}$ converges. By the Cauchy Criterion, given $\epsilon = \frac{1}{2}$. There exists N such that $\left| \sum_{k=n+1}^{2n} \frac{1}{k} \right| < \frac{1}{2}$ for all $m > n > N$. This contradicts to the above fact that $|S_{2n} - S_n| \geq \frac{1}{2}$, where m is chosen to be $2n$. \square

Note. The divergence of the harmonic series appears to have been established by Nicole Oresme (1323?-

1382) by showing that the sequence $\left\{ \sum_{k=1}^n \frac{1}{k} \right\}$ is NOT

bounded. The name *harmonic series* originated from the Greeks. Pythagoras studied the notes emitted by plucked strings of various lengths. If a string which emits middle C when plucked is reduced to two-thirds of its length, it will emit the note G (musicians call the interval from C to G a *fifth*). If the length of the string is halved, it will emit top C, i.e., an octave higher. These notes are fundamental to the Pythagorean theory of harmony, and the corresponding lengths

$$1, \frac{2}{3}, \frac{1}{2}$$

are said to be in *harmonic progression*. Note that their inverses form an arithmetic progression given by

$$1, \frac{3}{2}, 2.$$

The harmonic series defined above has similar properties, as inverses of the terms of the associated sequence

$$1, \frac{1}{2}, \frac{1}{3}, \dots$$

again forms the arithmetic progression $1, 2, 3, \dots$.

THEOREM 1.12. *Suppose that **eventually** $a_k \geq 0$. Then $\sum_{k=1}^{\infty} a_k$ converges **if and only if** $\{S_n\}$ is bounded above.*

PROOF. We may assume that $a_k \geq 0$ for all k . Since

$$S_{n+1} - S_n = a_{n+1} \geq 0,$$

the sequence $\{S_n\}$ is monotone increasing.

Thus $\sum_{k=1}^{\infty} a_k$ converges if and only if $\{S_n\}$ converges, if and only if $\{S_n\}$ is bounded above (by the Monotone Convergence Theorem). \square

Note. Suppose that **eventually** $a_k \geq 0$. This theorem means that the following.

1) If $\{S_n\}$ is bounded above, then $\sum_{k=1}^{\infty} a_k$ converges.

2) If $\{S_n\}$ is NOT bounded above, then $\sum_{k=1}^{\infty} a_k$ diverges.

2. Tests for Positive Series

A series $\sum_{k=1}^{\infty} a_k$ is called a **(eventually) positive series** if every term a_k is (eventually) positive.

2.1. Comparison Test.

THEOREM 2.1 (Comparison Test). *Consider two series $\sum_{k=1}^{\infty} a_k$ and $\sum_{k=1}^{\infty} b_k$. Suppose that eventually $0 \leq a_k \leq b_k$.*

(i) **If $\sum_{k=1}^{\infty} b_k$ converges, then $\sum_{k=1}^{\infty} a_k$ converges.**

(ii) **If $\sum_{k=1}^{\infty} a_k$ diverges, then $\sum_{k=1}^{\infty} b_k$ diverges.**

PROOF. Assertion (ii) follows immediately from (i).

We may assume that $0 \leq a_k \leq b_k$ for all k . Let $A_n = \sum_{k=1}^n a_k$, and $B_n = \sum_{k=1}^n b_k$. Then $A_n \leq B_n$ for all n .

Suppose that $\sum_{k=1}^{\infty} b_k$ converges, that is, $\sum_{k=1}^{\infty} b_k$ is a (finite) number. Then

$$A_n \leq B_n \leq \sum_{k=1}^{\infty} b_k$$

for all n and so A_n is bounded above.

By Theorem 1.12, $\sum_{k=1}^{\infty} a_k$ is convergent. □

EXAMPLE 2.2. The series $\sum_{k=1}^{\infty} \left(\frac{2k-1}{3k+2}\right)^k$ converges

because

$$\left(\frac{2k-1}{3k+2}\right)^k \leq \left(\frac{2}{3}\right)^k$$

and the geometric series $\sum_{k=1}^{\infty} \left(\frac{2}{3}\right)^k$ converges.

Remark. 1. Suppose $0 \leq a_n \leq b_n$ and $\sum_{n=1}^{\infty} b_n$ diverges. Then NO conclusion can be drawn.

2. Similarly, suppose $0 \leq a_n \leq b_n$ and $\sum_{n=1}^{\infty} a_n$ converges. Then NO conclusion can be drawn.

EXAMPLE 2.3.

$$0 \leq \left(\frac{1}{2}\right)^n \leq 2^n \leq 3^n.$$

The series $\sum_{n=1}^{\infty} 3^n$ diverges. Now $\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n$ converges

and $\sum_{n=1}^{\infty} 2^n$ diverges.

COROLLARY 2.4 (Limit Comparison Test). *Suppose that $\sum_{k=1}^{\infty} a_k$ and $\sum_{k=1}^{\infty} b_k$ are (eventually) positive series.*

(a). *If $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = L$ with $0 < L < \infty$, then $\sum_{k=1}^{\infty} a_k$*

converges if and only if $\sum_{k=1}^{\infty} b_k$ converges. (So

$\sum_{k=1}^{\infty} a_k$ diverges if and only if $\sum_{k=1}^{\infty} b_k$ diverges.)

(b). *If $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0$, (that is, $a_k \ll b_k$), and $\sum_{k=1}^{\infty} b_k$*

converges, then $\sum_{k=1}^{\infty} a_k$ converges.

(c). *If $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0$, (that is, $a_k \ll b_k$), and $\sum_{k=1}^{\infty} a_k$*

diverges, then $\sum_{k=1}^{\infty} b_k$ diverges.

Remark. If $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = +\infty$, interchange a_n and b_n , and then apply assertions (b) and (c).

PROOF. We may assume that $\sum_{k=1}^{\infty} a_k$ and $\sum_{k=1}^{\infty} b_k$ are positive series.

(a). Since

$$\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = L \quad (\neq 0, \neq \infty),$$

$\left\{ \frac{a_k}{b_k} \right\}$ is bounded above, say, by M . Thus

$$0 \leq a_k \leq M b_k$$

for all k . Similarly, since $\lim_{k \rightarrow \infty} \frac{b_k}{a_k} = \frac{1}{L} \quad (\neq 0, \neq \infty)$,

$\left\{ \frac{b_k}{a_k} \right\}$ is bounded above, say, by M' . Thus

$$0 \leq b_k \leq M' a_k$$

for all k .

If $\sum_{k=1}^{\infty} a_k$ is convergent, then $\sum_{k=1}^{\infty} M' a_k = M' \sum_{k=1}^{\infty} a_k$ is also convergent. By the comparison test, it follows that $\sum_{k=1}^{\infty} b_k$ is also convergent because $b_k \leq M' a_k$.

If $\sum_{k=1}^{\infty} b_k$ is convergent, then $\sum_{k=1}^{\infty} Mb_k = M \sum_{k=1}^{\infty} b_k$ is also convergent. By the comparison test, it follows that $\sum_{k=1}^{\infty} a_k$ is also convergent because $a_k \leq Mb_k$.

Hence $\sum_{k=1}^{\infty} a_k$ is convergent if and only if $\sum_{k=1}^{\infty} b_k$ is convergent. Hence $\sum_{k=1}^{\infty} a_k$ is divergent if and only if $\sum_{k=1}^{\infty} b_k$ is divergent.

Now we prove assertions (b) and (c).

$$\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0$$

implies for every $\epsilon > 0$, there is an N such that

$$\left| \frac{a_k}{b_k} - 0 \right| < \epsilon \quad \forall k > N.$$

We choose $\epsilon = 1$. Then the above inequality is

$$a_k < b_k \quad \forall k > N.$$

We get the result by applying the comparison test. \square

Standard series used in comparison and limit comparison tests.

1. The Geometric Series:

$$\sum_{n=1}^{\infty} ar^{n-1} = \begin{cases} \text{converges} & \text{if } |r| < 1, \\ \text{diverges} & \text{if } |r| \geq 1. \end{cases}$$

2. The p-series: for a fixed p ,

$$\sum_{n=1}^{\infty} \frac{1}{n^p} = \begin{cases} \text{converges} & \text{if } p > 1, \\ \text{diverges} & \text{if } p \leq 1. \end{cases}$$

To be proved in the subsection on Integral Test.

EXAMPLE 2.5. *Determine the convergence or divergence:*

$$1) \sum_{n=1}^{\infty} \frac{1 + \cos n}{n^2}$$

$$2) \sum_{n=1}^{\infty} \frac{\ln n + n^3 + 8}{n^4 - 2n + 3}$$

SOLUTION. (1). It is convergent, by the comparison test, because

$$0 \leq \frac{1 + \cos n}{n^2} \leq \frac{2}{n^2}$$

and the p -series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is convergent.

(2). It is divergent, by the limit comparison test, because

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\frac{\ln n + n^3 + 8}{n^4 - 2n + 3}}{\frac{1}{n}} &= \lim_{n \rightarrow \infty} \frac{n \ln n + n^4 + 8n}{n^4 - 2n + 3} \\ &= \lim_{n \rightarrow \infty} \frac{\frac{\ln n}{n^3} + 1 + \frac{1}{n^3}}{1 - \frac{2}{n^3} + \frac{3}{n^4}} = \frac{0 + 1 + 0}{1 - 0 + 0} = 1 \end{aligned}$$

and the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. □

EXAMPLE 2.6. *Determine convergence or divergence of $\sum_{n=2}^{\infty} \frac{1}{(\ln n)^k}$, where k is a constant.*

SOLUTION. Let $b_n = \frac{1}{(\ln n)^k}$ and let $a_n = \frac{1}{n}$. Then

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{\frac{1}{(\ln n)^k}} = \lim_{n \rightarrow \infty} \frac{(\ln n)^k}{n} = 0.$$

Since the harmonic series $\sum_{n=2}^{\infty} \frac{1}{n}$ is divergent, the series

$\sum_{n=2}^{\infty} \frac{1}{(\ln n)^k}$ is divergent for any k . □

2.2. Integral Test. Let $f(x)$ be a real-valued function on $[a, +\infty)$ such that $f(x)$ is Riemann integrable, that is, the integral $\int_a^b f(x) dx$ exists for every $b > a$. The following theorem is useful.

THEOREM 2.7. *Let $f(x)$ be a real-valued function on $[a, b]$.*

- (a). **If $f(x)$ is continuous on $[a, b]$, then $f(x)$ is Riemann integrable on $[a, b]$.**
- (b). **If $f(x)$ is monotone on $[a, b]$, then $f(x)$ is Riemann integrable on $[a, b]$.**

The improper integral is defined by

$$\begin{aligned}\int_a^\infty f(x) dx &= \lim_{b \rightarrow \infty} \int_a^b f(x) dx \\ &= \text{area under } f(x) \text{ over } [a, \infty).\end{aligned}$$

Here we say that $\int_a^\infty f(x) dx$ **converges** if the limit

$$\lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

exists (finite), i.e., the area under $f(x)$ over $[a, \infty)$ is finite.

We also say that $\int_a^\infty f(x) dx$ **diverges** if the limit


$$\lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

does not exist.

THEOREM 2.8 (Integral Test). *Let $f(x)$ be an (eventually) positive monotone decreasing function on $[1, +\infty)$. Suppose we have a series $\sum_{k=1}^{\infty} a_k$ such that $a_k = f(k)$, then the series $\sum_{k=1}^{\infty} a_k$ and the integral $\int_1^{\infty} f(x)dx$ either both converge or both diverge.*

PROOF. We may assume that $f(x)$ is positive monotone decreasing on $[1, +\infty)$.

Let $a_n = f(n)$ for all n .



From the graph, we see that

area of the rectangles \leq area under $f(x)$ over $[1, n]$, i.e.,

$$\sum_{k=2}^n f(k) \leq \int_1^n f(x) dx \leq \int_1^{\infty} f(x) dx.$$


Thus, if $\int_1^{\infty} f(x) dx < \infty$, then

$$\sum_{k=2}^n a_k = \sum_{k=2}^n f(k) \leq \int_1^{\infty} f(x) dx < \infty,$$

i.e., for all n , $\sum_{k=2}^n a_k$ is bounded above by the finite number $\int_1^{\infty} f(x) dx$.

Since we also have $a_k \geq 0$, it follows from Theorem 1.12 that $\sum_{k=2}^{\infty} a_k$ converges, and thus $\sum_{k=1}^{\infty} a_k$ also converges.

Next we consider the following graph:



From the graph, it is easy to see that

area under the rectangles \geq area under $f(x)$ over $[1, n]$,

i.e., $\sum_{k=1}^{n-1} f(k) \geq \int_1^n f(x) dx$. Thus, if $\sum_{k=1}^{\infty} a_k < \infty$,
then

$$\infty > \sum_{k=1}^{\infty} a_k \geq \sum_{k=1}^{n-1} a_k = \sum_{k=1}^{n-1} f(k) \geq \int_1^n f(x) dx,$$

i.e., $\int_1^n f(x) dx$ is bounded above by the finite number
 $\sum_{k=1}^{\infty} a_k < \infty$.

Letting $n \rightarrow \infty$, it follows that we have

$$\int_1^{\infty} f(x) dx \leq \sum_{k=1}^{\infty} a_k < \infty.$$

In conclusion, we have $\sum_{k=1}^{\infty} a_k$ converges if and only if $\int_1^{\infty} f(x) dx$ converges, which also means that $\sum_{k=1}^{\infty} a_k$ diverges if and only if $\int_1^{\infty} f(x) dx$ diverges. \square

EXAMPLE 2.9. *Show that*

1) *the series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.*

2) *the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^k}$ converges if and only if $k > 1$.*

PROOF. (1). If $p \leq 0$, then $\frac{1}{n^p}$ does not tend to 0 and so, by divergence test, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges.

Assume that $p > 0$. Let $f(x) = \frac{1}{x^p}$ on $[1, +\infty)$. Then $f(x)$ is positive monotone decreasing. Now

$$\int_1^{\infty} f(x) dx = \int_1^{\infty} x^{-p} dx = \begin{cases} \frac{1}{-p+1} x^{-p+1} \Big|_1^{+\infty} & p \neq 1 \\ \ln(+\infty) - \ln 1 & p = 1 \end{cases}$$

Thus $\int_1^{\infty} f(x) dx$ converges if and only if $p > 1$ and so $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

(2). the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^k}$ converges **if and only if**
 $k > 1$.

Let $f(x) = \frac{1}{x(\ln x)^k}$ on $[2, +\infty)$. Then $f(x)$ is positive. We check that $f(x)$ is *eventually* monotone decreasing. From

$$\begin{aligned} f'(x) &= (x^{-1}(\ln x)^{-k})' = -x^{-2}(\ln x)^{-k} - kx^{-1}(\ln x)^{-k-1} \frac{1}{x} \\ &= -x^{-2}(\ln x)^{-k-1}(\ln x + k), \end{aligned}$$

we have $f'(x) \leq 0$ when $\ln x > -k$. Thus $f(x)$ is monotone decreasing when $\ln x > -k$ and so $f(x)$ is eventually monotone decreasing.

Now

$$\begin{aligned} \int_2^{\infty} f(x) dx &= \int_2^{\infty} \frac{1}{x(\ln x)^k} dx \frac{y = \ln x}{dy = \frac{1}{x} dx} \int_{\ln 2}^{\infty} \frac{1}{y^k} dy \\ &= \begin{cases} \frac{1}{-k+1} y^{-k} \Big|_{\ln 2}^{\infty} & k \neq 1 \\ \ln(+\infty) - \ln(\ln 2) = +\infty & k = 1 \end{cases} \end{aligned}$$

Thus $\int_2^{\infty} f(x) dx$ converges if and only if $k > 1$ and so the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^k}$ converges if and only if $k > 1$. □

2.3. Ratio Test.

THEOREM 2.10 (Ratio Test). *Consider the positive series $\sum_{n=1}^{\infty} a_n$. Suppose*

$$(1) \quad \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \ell.$$

(i) *If $0 \leq \ell < 1$, then $\sum_{n=1}^{\infty} a_n$ converges.*

(ii) *If $1 < \ell \leq \infty$, then $\sum_{n=1}^{\infty} a_n$ diverges.*

(iii) *If $\ell = 1$, then the test is inconclusive.*

PROOF. We will prove (i) and (ii). Given any $\epsilon > 0$, it follows from (1) that there exists N such that for all $n > N$,

$$\left| \frac{a_{n+1}}{a_n} - \ell \right| < \epsilon \quad \text{or} \quad \ell - \epsilon < \frac{a_{n+1}}{a_n} < \ell + \epsilon.$$

By repeating using the above inequalities, it follows that for all $m > 0$,

$$(\ell - \epsilon)^m < \frac{a_{N+2}}{a_{N+1}} \cdot \frac{a_{N+3}}{a_{N+2}} \cdots \frac{a_{N+1+m}}{a_{N+m}} = \frac{a_{N+1+m}}{a_{N+1}} < (\ell + \epsilon)^m$$

$$(2) \quad a_{N+1}(\ell - \epsilon)^m < a_{N+1+m} < a_{N+1}(\ell + \epsilon)^m.$$

(i). If $\ell < 1$, choose $\epsilon > 0$ such that $\ell + \epsilon < 1$, then $\sum_{m=1}^{\infty} a_{N+1}(\ell + \epsilon)^m$ converges (since it is a geometric series with common ratio satisfying $|r| = \ell + \epsilon < 1$).

Together with the right-hand-side of (2), it follows from the comparison test that $\sum_{m=1}^{\infty} a_{N+1+m}$ converges,

and thus $\sum_{n=1}^{\infty} a_n$ converges.

(ii). If $\ell > 1$, choose $\epsilon > 0$ such that $\ell - \epsilon > 1$, then by the left-hand-side of (2), we have, for all $m > 0$,

$$a_{N+1+m} \geq a_{N+1}(\ell - \epsilon)^m > a_{N+1} > 0.$$

In particular, $\lim_{n \rightarrow \infty} a_n \neq 0$ or does not exist. By the divergence test, $\sum_{n=1}^{\infty} a_n$ diverges. \square

EXAMPLE 2.11. *Determine convergence or divergence.*

$$1) \sum_{n=1}^{\infty} \frac{n!}{n^n}.$$

$$2) \sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!}.$$

SOLUTION. (1).

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{\frac{(n+1)!}{(n+1)^{n+1}}}{\frac{n!}{n^n}} = \lim_{n \rightarrow \infty} \frac{(n+1)!}{n! \cdot \frac{(n+1)^n}{n^n} \cdot (n+1)} \\ &= \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^n} = \frac{1}{e} < 1. \end{aligned}$$

Thus the series converges.

(2).

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{\frac{[(n+1)!]^2}{(2n+2)!}}{\frac{(n!)^2}{(2n)!}} = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{(2n+2)(2n+1)} \\ &= \lim_{n \rightarrow \infty} \frac{(n+1)^2/n^2}{(2n+2)(2n+1)/n^2} = \lim_{n \rightarrow \infty} \frac{1 + 2/n + 1/n^2}{(2 + 2/n)(2 + 1/n)} \end{aligned}$$

$$= \frac{1 + 0 + 0}{(2 + 0) \cdot (2 + 0)} = \frac{1}{4} < 1.$$

Thus the series converges.



2.4. Root Test.

THEOREM 2.12 (Root Test). *Consider the series*
 $\sum_{n=1}^{\infty} a_n$ *with each*
 $a_n \geq 0$, *and let*

$$(3) \quad \ell = \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

(i) *If* $0 \leq \ell < 1$, *then* $\sum_{n=1}^{\infty} a_n$ **converges**.

(ii) *If* $1 < \ell \leq \infty$, *then* $\sum_{n=1}^{\infty} a_n$ **diverges**.

(iii) *If* $\ell = 1$, *then the test is* **inconclusive**.

PROOF. We will prove (i) and (ii).

(i). Suppose that $\ell < 1$. Then for all given $\epsilon > 0$, it follows from (3) and Proposition 8.9 of chapter 1, that there exists an N such that $\sqrt[n]{a_n} < \ell + \epsilon$ for all $n > N$. Now choose $\epsilon > 0$ s.t. $\ell + \epsilon < 1$. Then

$$(4) \quad 0 \leq a_n < (\ell + \epsilon)^n \quad \text{for all } n > N.$$

Since $\sum_{n=1}^{\infty} (\ell + \epsilon)^n$ converges (as it is a geometric series with common ratio satisfying $|r| = \ell + \epsilon < 1$), it follows from (4) and the comparison test that $\sum_{n=1}^{\infty} a_n$ converges.

(ii). We are going to prove (ii) by contradiction. Given that $\ell > 1$. Suppose that $\sum_{n=1}^{\infty} a_n$ converges. Then by Theorem 1.7, we have $\lim_{n \rightarrow \infty} a_n = 0$. In particular, there exists N such that $0 \leq a_n < 1$ for all $n > N$. Hence $\sqrt[n]{a_n} < 1$ for all $n > N$, and it follows that we must have $\ell \leq 1$, which is a contradiction. Hence $\sum_{n=1}^{\infty} a_n$ diverges. \square

COROLLARY 2.13 (Simplified Root Test). *Consider the series $\sum_{n=1}^{\infty} a_n$ with each $a_n \geq 0$. Suppose that $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \ell$.*

- (i) *If $0 \leq \ell < 1$, then $\sum_{n=1}^{\infty} a_n$ **converges**.*
- (ii) *If $1 < \ell \leq \infty$, then $\sum_{n=1}^{\infty} a_n$ **diverges**.*
- (iii) *If $\ell = 1$, then the test is **inconclusive**.*

PROOF. We will prove (i) and (ii). Recall from Theorem 8.13 of chapter 1 that if $\lim_{n \rightarrow \infty} \sqrt[n]{a_n}$ exists, then $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n}$. Then the Corollary follows from Theorem 2.12. □

EXAMPLE 2.14. *Determine convergence or divergence of the series*

$$\sum_{n=1}^{\infty} 2^n \left(1 - \frac{1}{n}\right)^{n^2}.$$

SOLUTION.

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{a_n} &= \lim_{n \rightarrow \infty} \left[2^n \left(1 - \frac{1}{n}\right)^{n^2} \right]^{\frac{1}{n}} \\ &= \lim_{n \rightarrow \infty} 2 \left(1 - \frac{1}{n}\right)^n = \frac{2}{e} < 1. \end{aligned}$$

Thus the series converges.

□

EXAMPLE 2.15. *Determine convergence or divergence of the series*

$$\sum_{n=1}^{\infty} (3 + \sin n)^n \left(1 - \frac{2}{n}\right)^{n^2}.$$

PROOF.

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{a_n} &= \overline{\lim}_{n \rightarrow \infty} \left[(3 + \sin n)^n \left(1 - \frac{2}{n}\right)^{n^2} \right]^{\frac{1}{n}} \\ &= \overline{\lim}_{n \rightarrow \infty} (3 + \sin n) \left(1 - \frac{2}{n}\right)^n \leq \overline{\lim}_{n \rightarrow \infty} 4 \left(1 - \frac{2}{n}\right)^n = \frac{4}{e^2} < 1.\end{aligned}$$

Thus the series converges. \square

3. Alternating Series

An **alternating series** is of the form

$$\sum_{n=1}^{\infty} (-1)^{n+1} a_n = a_1 - a_2 + a_3 - a_4 + \cdots, \quad \text{or}$$

$$\sum_{n=1}^{\infty} (-1)^n a_n = -a_1 + a_2 - a_3 + a_4 - \cdots$$

with each $a_n > 0$.

EXAMPLE 3.1.

$$\begin{aligned} &1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \cdots \\ &\quad 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots \\ &\quad -1 + 2 - 3 + 4 - 5 + 6 - \cdots \end{aligned}$$

THEOREM 3.2 (The Alternating Series test). *If $\{b_n\}$ is a sequence satisfying*

- (i) $b_1 \geq b_2 \geq b_3 \geq \cdots \geq 0$, and
- (ii) $\lim_{n \rightarrow \infty} b_n = 0$,

then $\sum_{n=1}^{\infty} (-1)^{n+1} b_n$ (and $\sum_{n=1}^{\infty} (-1)^n b_n$) converge.

Note. The test holds if $\{b_n\}$ is **eventually positive** and **eventually monotone decreasing** with $\lim_{n \rightarrow \infty} b_n = 0$.

PROOF. We prove that the sequence of partial sums $S_n = \sum_{k=1}^n (-1)^{k+1} b_k$ is Cauchy.

For $m > n$, note that

$$\begin{aligned} & |S_m - S_n| \\ &= |(-1)^{n+2} b_{n+1} + (-1)^{n+3} b_{n+2} + \cdots + (-1)^{m+1} b_m| \\ &= |b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots + (-1)^{m-n-1} b_m| \\ &\leq b_{n+1} + b_m \end{aligned}$$

because

$$\begin{aligned} & b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots + (-1)^{m-n-1} b_m \\ &= (b_{n+1} - b_{n+2}) + (b_{n+3} - b_{n+4}) + \cdots \pm b_m \geq -b_m \end{aligned}$$

and

$$\begin{aligned} & b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots + (-1)^{m-n-1} b_m \\ &= b_{n+1} - (b_{n+2}) - (b_{n+3}) - (b_{n+4} - b_{n+5}) + \cdots \leq b_{n+1} + b_m, \end{aligned}$$

where we use the property that $\{b_n\}$ is positive and monotone decreasing.

Given any $\epsilon > 0$, there exists N such that $b_n = |b_n - 0| < \frac{\epsilon}{2}$. Then for $m > n > N$,

$$|S_m - S_n| \leq b_{n+1} + b_m < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Thus $\{S_n\}$ is Cauchy and hence the result. □

EXAMPLE 3.3. *Show that convergence or divergence of the series*

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n^p} = \begin{cases} \text{convergence} & p > 0 \\ \text{divergence} & p \leq 0. \end{cases}$$

PROOF. If $p \leq 0$, then the n -th term $(-1)^n \frac{1}{n^p}$ does not tend to 0. Thus the series diverges in this case by the divergence test.

Assume that $p > 0$. Let $a_n = \frac{1}{n^p}$. Then $a_n > 0$, monotone decreasing and $\lim_{n \rightarrow \infty} a_n = 0$. Thus the alternating series converges in this case.

In conclusion, we have that the series $\sum_{n=1}^{\infty} (-1)^n \frac{1}{n^p}$ converges when $p > 0$ and diverges when $p \leq 0$. \square

Now we are going to give a theorem providing an estimate on the sum of (certain) series. For instance,

let $S = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^2}$. By taking the partial sum, we have

$$S \approx S_1 = 1, \quad S \approx S_2 = 1 - \frac{1}{2^2} = 0.75,$$

$$S \approx S_3 = 1 - \frac{1}{2^2} + \frac{1}{3^2} \approx 0.861, \dots$$

By using computer program, we are able to compute much more, say $S_{1000000}$. A mathematical problem is then what is the ‘error’ for estimating S by using the partial sum S_n . In other words, how to estimate the remainder

$$R_n = |S - S_n| = |a_{n+1} + a_{n+2} + \dots|.$$

THEOREM 3.4 (Alternating Series Estimation). *Let $\{b_n\}$ be a sequence satisfying*

(i) $b_1 \geq b_2 \geq b_3 \geq \dots \geq 0$, and

(ii) $\lim_{n \rightarrow \infty} b_n = 0$.

Let

$$S_n = \sum_{k=1}^n (-1)^{k+1} b_k \quad \text{and} \quad S = \sum_{k=1}^{\infty} (-1)^{k+1} b_k$$

Then the remainder $R_n = |S - S_n| \leq b_{n+1}$ for all n .

PROOF.

$$\begin{aligned} R_n &= \left| (-1)^{n+2}b_{n+1} + (-1)^{n+3}b_{n+2} + \cdots \right| \\ &= |b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots|. \end{aligned}$$

Since

$$\begin{aligned} & b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots \\ = & b_{n+1} - (b_{n+2} - b_{n+3}) - (b_{n+4} - b_{n+5}) - (b_{n+6} - b_{n+7}) - \cdots \\ & \leq b_{n+1} \end{aligned}$$

and

$$\begin{aligned} & b_{n+1} - b_{n+2} + b_{n+3} - b_{n+4} + \cdots \\ = & (b_{n+1} - b_{n+2}) + (b_{n+3} - b_{n+4}) + \cdots \geq 0, \end{aligned}$$

we have $R_n \leq a_{n+1}$. \square

EXAMPLE 3.5. Estimate $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^4}$ with error within 0.001.

SOLUTION. From $\frac{1}{(n+1)^4} \leq 10^{-3}$, we have $n+1 \geq 6$ or $n \geq 5$. Thus

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^4} \approx 1 - \frac{1}{2^4} + \frac{1}{3^4} - \frac{1}{4^4} + \frac{1}{5^4}$$

with error within 0.001. \square

4. Absolute and Conditional Convergence

DEFINITION 4.1. A series $\sum_{n=1}^{\infty} a_n$ is called **absolutely convergent** if $\sum_{n=1}^{\infty} |a_n|$ converges.

THEOREM 4.2. Every absolutely convergent series is convergent.

PROOF. Suppose that $\sum_{n=1}^{\infty} a_n$ converges absolutely,

that is, $\sum_{n=1}^{\infty} |a_n|$ converges by the definition. Let $T_n =$

$$\sum_{k=1}^n |a_k|, \quad S_n = \sum_{k=1}^n a_k.$$

Since $\{T_n\}$ converges, $\{T_n\}$ is Cauchy.

Thus, for any $\epsilon > 0$, there is a N such that $|T_n - T_m| < \epsilon$ for all $n, m > N$. For any $n, m > N$, we may assume that $m \geq n$, say $m = n + p$ (as one of them should be greater than another). Then

$$\begin{aligned} |S_n - S_m| &= |S_n - (S_n + a_{n+1} + a_{n+2} + \cdots + a_{n+p})| \\ &= |a_{n+1} + a_{n+2} + \cdots + a_{n+p}| \\ &\leq |a_{n+1}| + |a_{n+2}| + \cdots + |a_{n+p}| = T_m - T_n = |T_n - T_m| < \epsilon. \end{aligned}$$

Thus $\{S_n\}$ is a Cauchy sequence and so it converges.

Thus the series $\sum_{n=1}^{\infty} a_n$ converges and hence the result. □

EXAMPLE 4.3. *Determine convergence or divergence of the series*

$$\sum_{n=2}^{\infty} \frac{\sin n + \frac{1}{2}}{n(\ln n)^2}.$$

SOLUTION. Since

$$\left| \frac{\sin n + \frac{1}{2}}{n(\ln n)^2} \right| \leq \frac{|\sin n| + \frac{1}{2}}{n(\ln n)^2} \leq \frac{1 + \frac{1}{2}}{n(\ln n)^2}$$

and $\sum_{n=2}^{\infty} \frac{1 + \frac{1}{2}}{n(\ln n)^2} = \frac{3}{2} \sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$ converges by Exam-

ple 2.9, the series $\sum_{n=2}^{\infty} \left| \frac{\sin n + \frac{1}{2}}{n(\ln n)^2} \right|$ converges. Thus the

series $\sum_{n=2}^{\infty} \frac{\sin n + \frac{1}{2}}{n(\ln n)^2}$ converges. □

Remark. If you are testing for absolute convergence, all the techniques for the positive series are applicable.

Q: Is the converse of the Corollary true? I.e., if a series is convergent, will it be absolutely convergent?

A: No, it is not necessarily true.

EXAMPLE 4.4. The series $\sum_{n=1}^{\infty} (-1)^n \frac{1}{n}$ converges by Example 3.3, but it is NOT absolutely convergent by the p -series.

DEFINITION 4.5. A series $\sum_{n=1}^{\infty} a_n$ is said to be **conditionally convergent** if $\sum_{n=1}^{\infty} a_n$ converges but $\sum_{n=1}^{\infty} |a_n|$ diverges.

EXAMPLE 4.6. The series $\sum_{n=1}^{\infty} (-1)^n \frac{1}{\sqrt{n}}$ is conditionally convergent.

REMARK 4.7. Every series is either absolutely convergent, conditionally convergent or divergent. \square

EXAMPLE 4.8. The series

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n^p} = \begin{cases} \text{absolutely convergence} & p > 1 \\ \text{conditionally convergence} & 0 < p \leq 1 \\ \text{divergence} & p \leq 0 \end{cases}$$

5. Rearrangement properties of a series - an informal treatment

Roughly speaking, a **rearrangement** of a given series is another series produced from a given series by scrambling the order of the terms of the given series, but using all the terms exactly once. **Example:** Consider the harmonic series

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \cdots .$$

An rearrangement of the harmonic series is obtained by interchanging the 1st and 2nd terms, the 3rd and 4th terms, and so on, resulting in the new series

$$\frac{1}{2} + 1 + \frac{1}{4} + \frac{1}{3} + \frac{1}{6} + \frac{1}{5} + \frac{1}{8} + \frac{1}{7} + \cdots .$$

It turns out that the sum of a rearrangement of a given series may be different from the original series. In fact, for any given **conditional convergent** series

$\sum_{n=1}^{\infty} a_n$ and any given real number $c \in \mathbb{R}$, it is known

that one may rearrange $\sum_{n=1}^{\infty} a_n$ so that the new series

$\sum_{n=1}^{\infty} b_n$ will converge to c (i.e., $\sum_{n=1}^{\infty} b_n = c$). On the

other hand, an **absolutely convergent** series has

much nicer rearrangement properties. Given an absolutely convergent series $\sum_{n=1}^{\infty} a_n$, whose sum is denoted

by S , i.e., $\sum_{n=1}^{\infty} a_n = S$. Then for any rearrangement

$\sum_{n=1}^{\infty} b_n$ of $\sum_{n=1}^{\infty} a_n$, one always has $\sum_{n=1}^{\infty} b_n = S$. In other

words, rearranging an absolutely convergent series will not change the sum.

Reading Exercise: See Bartle-Sherbert[1, p.255] for details.

6. Remarks on the various tests for convergence/divergence of series

1. n -th term test for divergence:

- a test for divergence ONLY, and it works for series with positive and negative terms, e.g. $\sum_{n=1}^{\infty} (-1)^n$.

2. Comparison test/Limit Comparison test:

- when applying these tests, one usually compares the given series with a geometric series or a p -series.

- generally works for series which look like the geometric series or the p -series,

e.g. $\sum_{n=1}^{\infty} \frac{2 + (-1)^n}{4^n}$, $\sum_{n=1}^{\infty} \frac{2^{\frac{1}{n}}}{n^2}$.

- when an oscillating factor/term appears, e.g. $\sum_{n=1}^{\infty} \frac{2 + (-1)^n}{3^n}$,
try the Comparison test rather than the Limit Comparison test.

3. Integral test:

e.g. $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$.

4. **Ratio test:**

- generally works for series which look like the geometric series, series with $n!$, and certain series defined recursively,

$$\text{e.g. } \sum_{n=1}^{\infty} \frac{n^2}{3^n}, \quad \sum_{n=1}^{\infty} \frac{(2n)!}{4^n \cdot n!},$$

$$\sum_{n=1}^{\infty} a_n, \text{ where } a_1 = 1, a_n = \left(\frac{1}{2} + \frac{1}{n}\right)a_{n-1}, n = 2, 3, \dots$$

5. **(Simplified) Root test:**

- generally works for series where a_n involves a high power such as the n -th power,

$$\text{e.g. } \sum_{n=1}^{\infty} \frac{n}{3^n}, \quad \sum_{n=1}^{\infty} 2^n \left(1 - \frac{1}{n}\right)^{n^2}.$$

6. **Alternating Series test:** - works for alternating series only,

$$\text{e.g. } \sum_{n=2}^{\infty} (-1)^n \frac{\ln n}{n}.$$

Remark. In general, Tests 2 - 5 works only for $\sum_{n=1}^{\infty} a_n$, where $a_n \geq 0$ (except Ratio and Root Tests for general series).

CHAPTER 3


Sequences and Series of Functions

1. Pointwise Convergence

1.1. Sequences of Functions. Let I be a (nonempty) subset of \mathbb{R} , e.g. $(-1, 1)$, $[0, 1]$, etc. For each $n \in \mathbb{N}$, let $F_n : I \rightarrow \mathbb{R}$ be a function. Then we say $\{F_n\}$ forms a **sequence of functions** on I .

EXAMPLE 1.1. (1). $F_n(x) = x^n$, $0 < x < 1$.
Then $\{F_n\}$ forms a sequence of functions on $(0,1)$.

$$\begin{aligned} F_1(x) &= x, & 0 < x < 1; \\ F_2(x) &= x^2, & 0 < x < 1; \\ F_3(x) &= x^3, & 0 < x < 1; \\ & \dots \end{aligned}$$



(2). $\left\{ \left(1 + \frac{x}{n}\right)^n \right\}$ forms a sequence of functions on $(-\infty, \infty)$. Write out some terms:


$$F_1(x) = 1+x \quad F_2(x) = \left(1 + \frac{x}{2}\right)^2 \quad F_3(x) = \left(1 + \frac{x}{3}\right)^3$$

If we fix the x , and let $n \rightarrow \infty$,

$$\lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x.$$

So for each x , we can define

$$F(x) = \lim_{n \rightarrow \infty} F_n(x).$$



DEFINITION 1.2. A sequence $\{F_n\}$ is said to **converge pointwise** to a function F on I if


$$\lim_{n \rightarrow \infty} F_n(x) = F(x) \quad \text{for each } x \in I,$$

i.e., for each $x \in I$ and given any $\epsilon > 0$, there exists an N (which depends on x and ϵ) such that

$$|F_n(x) - F(x)| < \epsilon \quad \forall n > N.$$

The function F is called the **limiting function** of $\{F_n\}$.

Graphically, this means



Remark. The limiting function is necessarily unique.

EXAMPLE 1.3. The sequence $\{x^n\}$ converges pointwise on $I = (0, 1)$ because the limit $F(x) = \lim_{n \rightarrow \infty} x^n = 0$ exists for any $0 < x < 1$.

The sequence $\{x^n\}$ does NOT converge on $[-1, 1]$ because the limit $\lim_{n \rightarrow \infty} x^n$ does not exist when $x = -1 \in [-1, 1]$.

1.2. Series of Functions. A series of functions on a set I is of the form

$$\sum_{n=1}^{\infty} f_n(x) = f_1(x) + f_2(x) + \cdots,$$

where each f_n is a function on I .

EXAMPLE 1.4. Below are some examples.

1. $\sum_{n=1}^{\infty} x^{n-1} = 1 + x + x^2 + x^3 + \cdots$
2. $\sum_{k=1}^{\infty} \frac{\sin kx}{k+x} = \frac{\sin x}{1+x} + \frac{\sin 2x}{2+x} + \frac{\sin 3x}{3+x} + \cdots,$ $0 \leq x \leq 1.$

As in chapter 2, we may form the **partial sums**

$$S_n(x) = \sum_{k=1}^n f_k(x) = f_1(x) + f_2(x) + \cdots + f_n(x).$$

Then $\{S_n\}$ forms a sequence of functions on I .

DEFINITION 1.5. The series $\sum_{n=1}^{\infty} f_n$ is said to **converge pointwise** (to a function S) on I if $\{S_n\}$ converges pointwise (to S) on I , (i.e. $\lim_{n \rightarrow \infty} S_n(x) = S(x)$ for each $x \in I$.)

EXAMPLE 1.6. What is the pointwise limit of $\sum_{n=1}^{\infty} x^{n-1}$,

where $x \in (-1, 1)$? Does $\sum_{n=1}^{\infty} x^{n-1}$ converge pointwise on $[-1, 1)$

SOLUTION. Consider the partial sum

$$S_n(x) = \sum_{i=1}^n x^{i-1} = 1 + x + \cdots + x^{n-1} = \frac{1 - x^n}{1 - x}$$

for $-1 < x < 1$. Thus

$$\sum_{n=1}^{\infty} x^{n-1} = \lim_{n \rightarrow \infty} S_n(x) = \lim_{n \rightarrow \infty} \frac{1 - x^n}{1 - x} = \frac{1}{1 - x}$$

for $x \in (-1, 1)$.

Since the series $\sum_{n=1}^{\infty} x^{n-1}$ diverges when $x = -1$, the series of functions $\sum_{n=1}^{\infty} x^{n-1}$ does NOT converge pointwise on $[-1, 1)$. \square

1.3. Some Questions on Pointwise Convergence. Suppose a sequence of functions $\{F_n\}$ converges pointwise to a function F on the interval $[a, b]$.

Also suppose a series of functions $\sum_{n=1}^{\infty} f_n(x)$ converges pointwise to a function $S(x)$ on $[a, b]$. Among the questions we want to consider the following. Some of these questions were incorrectly believed to be true by many mathematicians prior to nineteenth century, including the famous Cauchy. Cauchy in his text *Cours d'Analyse* “proved” a theorem to the effect that the limit of a convergent sequence of continuous functions was again continuous. As we will see, this result is false!

Question (a). If each F_n is continuous at $p \in [a, b]$, is F necessarily continuous at p ? Recall that F is continuous at p if and only if

$$\lim_{t \rightarrow p} F(t) = F(p).$$

Since $F(x) = \lim_{n \rightarrow \infty} F_n(x)$ for every $x \in [a, b]$, what we really asking is does

$$\lim_{t \rightarrow p} \left(\lim_{n \rightarrow \infty} F_n(t) \right) = \lim_{n \rightarrow \infty} \left(\lim_{t \rightarrow p} F_n(t) \right) ?$$

Question (a’). For series of functions, we can ask similar question. If each $f_n(x)$ is continuous at p , is

$S(x) = \sum_{n=1}^{\infty} f_n(x)$ necessarily continuous at p ? Again
what we really asking is does

$$\lim_{t \rightarrow p} \sum_{n=1}^{\infty} f_n(t) = \sum_{n=1}^{\infty} \lim_{t \rightarrow p} f_n(t) ?$$

Question (b). If each F_n is differentiable on $[a, b]$, is F necessarily differentiable on $[a, b]$? If so, does

$$\frac{d}{dx} \lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} \frac{d}{dx} F_n(x) ?$$

Question (b'). If each f_n is differentiable on $[a, b]$, is $S(x) = \sum_{n=1}^{\infty} f_n(x)$ necessarily differentiable on $[a, b]$? If so, does

$$\frac{d}{dx} \sum_{n=1}^{\infty} f_n(x) = \sum_{n=1}^{\infty} \frac{d}{dx} f_n(x) ?$$

Question (c). If each F_n is Riemann integrable on $[a, b]$, is F necessarily Riemann integrable on $[a, b]$? If so, does

$$\int_a^b \lim_{n \rightarrow \infty} F_n(x) dx = \lim_{n \rightarrow \infty} \int_a^b F_n(x) dx ?$$

Question (c'). If each f_n is Riemann integrable on $[a, b]$, is $S(x) = \sum_{n=1}^{\infty} f_n(x)$ necessarily Riemann integrable on $[a, b]$? If so, does

$$\int_a^b \sum_{n=1}^{\infty} f_n(x) dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx ?$$

Without **additional hypothesis** the answer to all of these questions is generally **no**. This additional hypothesis is so-called **uniform convergence**, introduced by Weierstrass in 1850. For his many contributions to the subject area, Weierstrass is often referred to as the father of modern analysis. The subsequent study on the subject together with questions from geometry and physics also lead to a new area called **topology** in the end of 19th century. Poincaré is often referred to as the father of topology. Topology together with Riemann geometry provide the mathematics foundation for Einstein's relativity theory. You may learn some basic knowledge of topology and modern geometry in 4000 and 5000 modules.

Below we only give counter-examples to Questions (a) and (c). You may read the text book [7] for more examples.

EXAMPLE 1.7 (Counter-example to Question (a)). Consider the functions

$$F_n(x) = x^n, \quad x \in [0, 1].$$

For each fixed $x \in [0, 1)$, we have

$$\lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} x^n = 0 \quad (\text{since } |x| < 1).$$

At $x = 1$, we have


$$\lim_{n \rightarrow \infty} \lim_{x \rightarrow 1} F_n(x) = \lim_{n \rightarrow \infty} F_n(1) = \lim_{n \rightarrow \infty} 1^n = 1.$$

Thus $\{F_n\}$ converges pointwise to the function F on the interval $[0, 1]$ given by

$$F(x) = \begin{cases} 0, & \text{for } x \in [0, 1), \\ 1, & x = 1. \end{cases}$$

Each F_n is continuous on the whole interval $[0, 1]$, but F is not continuous at $x = 1$. And


$$\lim_{x \rightarrow 1} \lim_{n \rightarrow \infty} F_n(x) = \lim_{x \rightarrow 1} F(x) = 0 \neq \lim_{n \rightarrow \infty} \lim_{x \rightarrow 1} F_n(x).$$



This simple example gives a counter example to Cauchy's (false) statement that the limit of continuous functions is continuous, namely the limit of continuous functions need not be continuous.

EXAMPLE 1.8 (Counter-example to Question (c)). Consider the functions

$$F_n(x) = \begin{cases} n^2 x, & 0 < x < \frac{1}{n}, \\ 2n - n^2 x, & \frac{1}{n} \leq x < \frac{2}{n}, \\ 0, & \frac{2}{n} \leq x < 1. \end{cases}$$



For each fixed $x \in (0, 1]$, one sees that $F_n(x) = 0$ whenever $n \geq \frac{2}{x}$, and hence

$$\lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} 0 = 0.$$

Also, at $x = 0$, we have

$$\lim_{n \rightarrow \infty} F_n(0) = \lim_{n \rightarrow \infty} n^2 \cdot 0 = 0.$$

Thus, $\{F_n\}$ converges pointwise to the zero function $F(x) \equiv 0$ on the interval $[0, 1]$.

For each $n \geq 1$, we have

$$\begin{aligned} \int_0^1 F_n(x) dx &= \int_0^{1/n} n^2 x dx + \int_{1/n}^{2/n} (2n - n^2 x) dx + \int_{2/n}^1 0 dx \\ &= \frac{n^2 x^2}{2} \Big|_0^{1/n} + \left(2nx - \frac{n^2 x^2}{2} \right) \Big|_{1/n}^{2/n} + 0 \\ &= \frac{1}{2} + (4 - 2) - \left(2 - \frac{1}{2} \right) + 0 = 1. \end{aligned}$$

Thus we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_0^1 F_n(x) dx &= \lim_{n \rightarrow \infty} 1 = 1 \neq 0 = \int_0^1 F(x) dx, \quad \text{i.e.} \\ \lim_{n \rightarrow \infty} \int_0^1 F_n(x) dx &\neq \int_0^1 \left(\lim_{n \rightarrow \infty} F_n(x) \right) dx. \end{aligned}$$

Reason: At different x , $F_n(x)$ converges to $F(x)$ at different pace (more specifically, in the definition of pointwise convergence, the choice of N depends on both ϵ and x).

2. Uniform Convergence

We define a slightly different concept of convergence.

2.1. Uniform Convergence of Sequences of Functions.

DEFINITION 2.1. $\{F_n\}$ is said to **converge uniformly** to a function F on a set I if for every $\epsilon > 0$, there exists an N (which depends only on ϵ) such that

$$|F_n(x) - F(x)| < \epsilon$$


for **ALL** $x \in I$ whenever $n > N$.

REMARK 2.2. *If $\{F_n\}$ converges uniformly to F on I , then $\{F_n\}$ converges pointwise to F on I . Conversely, if $\{F_n\}$ converges pointwise to F on I , then $\{F_n\}$ need not converge uniformly to F on I .*


The inequality in the definition can be expressed as

$$F(x) - \epsilon < F_n(x) < F(x) + \epsilon$$

for all $x \in I$ and $n > N$. The geometric interpretation is that for $n > N$ the graph of $y = F_n(x)$ lies in the band spanned by the curves $y = F(x) - \epsilon$ and $y = F(x) + \epsilon$, that is, given any $\epsilon > 0$ the graph of $y = F_n(x)$ **eventually** lies in the band spanned by the curves $y = F(x) - \epsilon$ and $y = F(x) + \epsilon$.



EXAMPLE 2.3. The sequence of function $F_n = x^n$ converges pointwise to 0 on $[0, 1)$.



But from the graph we can see that the graph of $F_n = x^n$ DOES NOT eventually lie in the band spanned by $y = -\epsilon$ and $y = +\epsilon$. The geometric reason also tells us that $\{x^n\}$ does not converge uniformly on $[0, 1)$.

2.2. Two Criteria for Uniform Convergence of $\{F_n\}$. The following theorem is useful (computationally) in determining whether a sequence of functions converges uniformly or not.

THEOREM 2.4 (*T*-test). *Suppose $\{F_n\}$ is a sequence of functions converging pointwise to a function F on a set I , and let*

$$T_n = \sup_{x \in I} |F_n(x) - F(x)|.$$

Then $\{F_n\}$ converges uniformly to F on I if and only if $\lim_{n \rightarrow \infty} T_n = 0$.

PROOF. First we prove the ‘only if’ part. Suppose that $\{F_n\}$ converges uniformly to F on I . Then for any given $\epsilon > 0$, there exists N such that

$$|F_n(x) - F(x)| < \frac{\epsilon}{2} \quad \text{for all } n > N \text{ and } x \in I$$

$$\implies T_n = \sup_{x \in I} |F_n(x) - F(x)| \leq \frac{\epsilon}{2} < \epsilon \quad \text{for all } n > N$$

$$\implies |T_n - 0| = T_n < \epsilon \quad \text{for all } n > N.$$

Hence we have $\lim_{n \rightarrow \infty} T_n = 0$.

Next we prove the ‘if’ part. Suppose that $\lim_{n \rightarrow \infty} T_n = 0$.

Then for any given $\epsilon > 0$, there exists N such that

$$|T_n - 0| = T_n < \epsilon \quad \text{for all } n > N$$

$$\implies \sup_{x \in I} |F_n(x) - F(x)| < \epsilon \quad \text{for all } n > N$$

$$\implies |F_n(x) - F(x)| < \epsilon \quad \text{for all } n > N \text{ and } x \in I.$$

Hence $\{F_n\}$ converges uniformly to F on I . This finishes the proof of the theorem. \square

THEOREM 2.5 (Cauchy's Criterion). *A sequence of functions $\{F_n\}$ converges uniformly on a set I if and only if given any $\epsilon > 0$, there exists a natural number N such that*

$$(5) \quad |F_n(x) - F_m(x)| < \epsilon \quad \text{for all } x \in I \text{ and all } m, n > N.$$

Remark: Here N does not depend on x .

PROOF. First we prove the ‘only if’ part. Suppose that $\{F_n\}$ converges uniformly to the function F on I .

Then given any $\epsilon > 0$, there exists N such that

$$|F_n(x) - F(x)| < \frac{\epsilon}{2} \quad \text{for all } x \in I \text{ and all } n > N.$$

Then for all $x \in I$ and $m, n > N$,

$$\begin{aligned} |F_n(x) - F_m(x)| &= |(F_n(x) - F(x)) - (F_m(x) - F(x))| \\ &\leq |F_n(x) - F(x)| + |F_m(x) - F(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

This finishes the proof of the ‘only if’ part.

Next we prove the ‘if’ part. Suppose that equation 5 holds.

Then for each fixed point $x \in I$, $\{F_n(x)\}$ is a Cauchy sequence of real numbers, and thus by Cauchy’s criterion for sequences, the sequence of real numbers $\{F_n(x)\}$ converges.

For each $x \in I$, we denote the limit by $F(x) = \lim_{n \rightarrow \infty} F_n(x)$. Then $\{F(x)\}_{x \in I}$ forms a function on I , which we denote by F .

Given any $\epsilon > 0$, by equation 5, there exists N such that

$$|F_n(x) - F_m(x)| < \frac{\epsilon}{2} \quad \text{for all } x \in I \text{ and all } m, n > N.$$

Then for each fixed $x \in I$ and $n > N$, we have

$$\begin{aligned} |F_n(x) - F(x)| &= |F_n(x) - \lim_{m \rightarrow \infty} F_m(x)| \\ &= \lim_{m \rightarrow \infty} |F_n(x) - F_m(x)| \leq \lim_{m \rightarrow \infty} \frac{\epsilon}{2} = \frac{\epsilon}{2} < \epsilon. \end{aligned}$$

Thus $\{F_n\}$ converges uniformly to F , and this finishes the proof of the ‘if’ part. \square

2.3. Examples.

EXAMPLE 2.6. Show that $F_n(x) = \frac{\sin^2 x}{n}$, $x \in (-\infty, +\infty)$, converges uniformly.

PROOF. The limiting function $F(x)$ is

$$F(x) = \lim_{n \rightarrow \infty} \frac{\sin^2 x}{n} = 0$$

for all $x \in (-\infty, +\infty)$. Since

$$\begin{aligned} T_n &= \sup_{x \in (-\infty, +\infty)} |F_n(x) - F(x)| \\ &= \sup_{x \in (-\infty, +\infty)} \left| \frac{\sin^2 x}{n} \right| \leq \frac{1}{n} \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, thus the sequence of functions $\{F_n(x)\}$ converges uniformly on $(-\infty, +\infty)$. \square

EXAMPLE 2.7. *Determine whether the following sequences of functions converge uniformly on the indicated interval.*

$$(a) f_n(x) = \frac{n^2 \ln x}{x^n}, x \in [1, +\infty);$$

$$(b) f_n(x) = \frac{n^2 \ln x}{x^n}, x \in [2, +\infty).$$

SOLUTION. Let $f(x) = \lim_{n \rightarrow \infty} f_n(x) = 0$ for $x \geq 1$.

$$(a). \quad T_n = \sup_{x \geq 1} |f_n(x) - 0| = \sup_{x \geq 1} \frac{n^2 \ln x}{x^n} = \sup_{x \geq 1} f_n(x).$$

From

$$f'_n(x) = n^2 \frac{1}{x} \cdot x^{-n} - n^3 \ln x \cdot x^{-n-1} = \frac{n^2 - n^3 \ln x}{x^{n+1}} = 0,$$

we have $n^2 - n^3 \ln x = 0$ or $x = e^{\frac{1}{n}}$. Observe that $f_n(x)$ is monotone increasing for $1 \leq x \leq e^{\frac{1}{n}}$ and monotone decreasing for $x \geq e^{\frac{1}{n}}$. Thus

$$T_n = \max_{x \geq 1} f_n(x) = f_n(e^{\frac{1}{n}}) = \frac{n^2 \cdot \frac{1}{n}}{\left(e^{\frac{1}{n}}\right)^n} = \frac{n}{e} \not\rightarrow 0$$

as $n \rightarrow \infty$ and so $\{f_n(x)\}$ does NOT converge uniformly.

(b). Since $e^{\frac{1}{n}} \leq 2$ for $n \geq 2$, the function $f_n(x)$ is monotone decreasing on $[2, +\infty)$ for $n \geq 2$ and so

$$T_n = \sup_{x \geq 2} |f_n(x) - f(x)| = f_n(2) = \frac{n^2 \ln 2}{2^n} \text{ for } n \geq 2.$$

Since $\lim_{n \rightarrow \infty} T_n = 0$, $\{f_n\}$ converges uniformly on $[2, +\infty)$. \square

EXAMPLE 2.8. Show that $F_n(x) = \frac{n^2 \ln x \sin nx}{x^n}$ converges uniformly on $[2, +\infty)$.

SOLUTION. $F(x) = \lim_{n \rightarrow \infty} F_n(x) = 0$ for $x \geq 2$. Observe

$$T_n = \sup_{x \geq 2} |F_n(x) - F(x)| = \sup_{x \geq 2} \frac{n^2 \ln x |\sin nx|}{x^n} \leq \frac{n^2 \ln 2}{2^n}$$

for $n \geq 2$. Since $\lim_{n \rightarrow \infty} T_n = 0$, $\{F_n\}$ converges uniformly. \square

Remark. Let $\{F_n\}$ be a sequence of functions on an interval I . To see whether $\{F_n\}$ is uniformly convergent, we may try to do by the following steps.

- (1). Determine the limiting function $F(x) = \lim_{n \rightarrow \infty} F_n(x)$.
- (2). Determine $T_n = \sup_{x \in I} |F_n(x) - F(x)|$.
- (3). Check whether $\lim_{n \rightarrow \infty} T_n = 0$.

If T_n is difficult to be determined, then we may try to estimate an upper bound of T_n (a lower bound of T_n if we guess that the sequence of functions might not be uniformly convergent).

2.4. Uniform Convergence of Series of Functions.

DEFINITION 2.9. $\sum_{n=1}^{\infty} f_n$ is said to **converge uniformly** (to S) on I if the sequence of its partial sums $\{S_n\}$ converges uniformly (to S) on I .

THEOREM 2.10 (*T*-test for Series of Functions).
 Suppose $\sum_{n=1}^{\infty} f_n(x)$ is a series of functions converging pointwise on a set I , and let

$$T_n = \sup_{x \in I} \left| \sum_{k=n+1}^{\infty} f_k(x) \right|.$$

Then $\sum_{n=1}^{\infty} f_n(x)$ **converges uniformly on I if and only if** $\lim_{n \rightarrow \infty} T_n = 0$.

PROOF. Let $S_n(x) = \sum_{k=1}^n f_k(x)$ and $S(x) = \sum_{k=1}^{\infty} f_k(x)$.

Then

$$|S_n(x) - S(x)| = \left| \sum_{k=n+1}^{\infty} f_k(x) \right|$$

and so the *T*-test, Theorem 2.4, applies. □

THEOREM 2.11 (Cauchy Criterion). *A series of functions $\sum_{n=1}^{\infty} f_n$ converges uniformly on a set I if and only if given any $\epsilon > 0$, there exists a natural number N such that*

$$\left| \sum_{k=n+1}^m f_k(x) \right| < \epsilon \quad \text{for all } x \in I \text{ and all } m > n > N.$$

Remark: Here N does not depend on x .

PROOF. The proof follows by applying the Cauchy Criterion to the partial sums $S_n(x) = \sum_{k=1}^n f_k(x)$. \square

The following test is very useful in verifying that certain series of functions converge uniformly to some functions on an interval.

THEOREM 2.12 (Weierstrass M -test). *Consider a series of functions $\sum_{k=1}^{\infty} f_k$ on a set I . Suppose that*

(i) $|f_k(x)| \leq M_k$ for all $x \in I$, $k = 1, 2, \dots$,

and

(ii) $\sum_{k=1}^{\infty} M_k$ converges.

Then $\sum_{k=1}^{\infty} f_k$ converges uniformly (to some function) on I .

Remark. Weierstrass M -test **only** states that if a series of functions satisfies conditions (i) and (ii), it converges uniformly on I . If a series of functions **does not** satisfy these two conditions, the test fails, namely, **no conclusion** that you can claim from this test.

PROOF. Since $\sum_{k=1}^{\infty} M_k$ converges (by (ii)), by the Cauchy Criterion, given any $\epsilon > 0$, there exists N such that

$$\sum_{k=n+1}^m M_k = \left| \sum_{k=n+1}^m M_k \right| < \epsilon \quad \text{for all } m > n > N$$

because $M_k \geq 0$.

Then for all $x \in I$, we have

$$\left| \sum_{k=n+1}^m f_k(x) \right| \leq \sum_{k=n+1}^m |f_k(x)| \leq \sum_{k=n+1}^m M_k < \epsilon.$$

Thus, by the Cauchy criterion, $\sum_{k=1}^{\infty} f_k$ converges uniformly on I . □

2.5. Examples.

EXAMPLE 2.13. Show that $\sum_{n=1}^{\infty} \frac{\cos^n x}{n^2 + x}$ converges uniformly on $(0, \infty)$.

PROOF. Since

$$\left| \frac{\cos^n x}{n^2 + x} \right| \leq \frac{1}{n^2}$$

for all $x \in (0, \infty)$ and the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is convergent

by the p -series, the series of functions $\sum_{n=1}^{\infty} \frac{\cos^n x}{n^2 + x}$ converges uniformly by the Weierstrass M -test and hence the result. \square

EXAMPLE 2.14. *Does the series of functions*

$$\sum_{n=1}^{\infty} n^2 x^n \sin nx$$

converge uniformly on the interval $[0, \frac{1}{2}]$? Justify your answer.

SOLUTION. Note that

$$\left| n^2 x^n \sin nx \right| \leq \frac{n^2}{2^n}$$

for $x \in [0, \frac{1}{2}]$. Since

$$\lim_{n \rightarrow \infty} \frac{\frac{(n+1)^2}{2^{n+1}}}{\frac{n^2}{2^n}} = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{2n^2} = \frac{1}{2} < 1,$$

the series $\sum_{n=1}^{\infty} \frac{n^2}{2^n}$ converges, by the ratio test,

and so the series of functions

$$\sum_{n=1}^{\infty} n^2 x^n \sin nx$$

converges uniformly on $[0, \frac{1}{2}]$ by the Weierstrass M -test. \square

EXAMPLE 2.15. Show that the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}$ converges uniformly on $[0, 1]$.

Note. The M -test fails for this example because

$$\sup_{x \in [0,1]} \left| (-1)^{n+1} \frac{x^n}{n} \right| = \frac{1}{n}$$

and $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. In this case, we use T -test or Cauchy Criterion.

PROOF. Let $a_n(x) = \frac{x^n}{n}$. Then, for $0 \leq x \leq 1$, we have

$$a_1(x) \geq a_2(x) \geq \cdots \geq 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n(x) = 0.$$

Thus the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}$ converges pointwise on $[0, 1]$ by the Alternating Series Test.

By the Alternating Series Estimation,

$$T_n = \sup_{0 \leq x \leq 1} \left| \sum_{k=n+1}^{\infty} (-1)^{k+1} \frac{x^k}{k} \right| \leq \sup_{0 \leq x \leq 1} \frac{x^{n+1}}{n+1} = \frac{1}{n+1}.$$

Since $\lim_{n \rightarrow \infty} \frac{1}{n+1} = 0$, the series $(-1)^{n+1} \frac{x^n}{n}$ converges uniformly by the T -test. \square

EXAMPLE 2.16. Show that the series $\sum_{n=1}^{\infty} \frac{x^n}{n}$ does not converge uniformly on $[0, 1)$.

PROOF. Observe that

$$\begin{aligned}
 T_n &= \sup_{0 \leq x < 1} |S_n(x) - S(x)| = \sup_{0 \leq x < 1} \left| \sum_{k=n+1}^{\infty} \frac{x^k}{k} \right| \\
 &= \sup_{0 < x < 1} \sum_{k=n+1}^{\infty} \frac{x^k}{k}. \\
 &= \sup_{0 < x < 1} \left(\frac{x^{n+1}}{n+1} + \frac{x^{n+2}}{n+2} + \dots \right) \\
 &\geq \sup_{0 < x < 1} \left(\frac{x^{n+1}}{n+1} + \frac{x^{n+2}}{n+2} + \dots + \frac{x^{2n}}{2n} \right) \\
 &= \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} \\
 &\quad \text{(because } \frac{x^k}{k} \text{ monotone increasing on } (0, 1)) \\
 &\geq \frac{1}{2n} + \frac{1}{2n} + \dots + \frac{1}{2n} = n \cdot \frac{1}{2n} = \frac{1}{2}.
 \end{aligned}$$

Thus the sequence $\{T_n\}$ does not tend to 0 and so the series of functions $\sum_{k=1}^{\infty} \frac{x^k}{k}$ does NOT converge uniformly on $[0, 1)$ by the T -test. \square

3. Uniform Convergence of $\{F_n\}$ and Continuity

In this section we will prove that the limit of uniformly convergent sequence of continuous functions is again continuous.

THEOREM 3.1. *Let $\{F_n\}$ be a sequence of continuous functions on an interval I . Suppose that $\{F_n\}$ converges uniformly to a function F on I . Then F is continuous on I .*

PROOF. Fix any point $x_0 \in I$. We are going to show that F is continuous at the point x_0 .

Given any $\epsilon > 0$, since $\{F_n\}$ converges uniformly to F on I , there exists N such that

$$|F_n(x) - F(x)| < \frac{\epsilon}{3} \quad \text{for all } x \in I \text{ and all } n > N.$$

Next we fix an $n > N$ (say, $n = [N] + 1$). Since F_n is continuous at x_0 , there exists $\delta > 0$ (here δ depends on x_0 and ϵ) such that for all x satisfying $|x - x_0| < \delta$, we have


$$|F_n(x) - F_n(x_0)| < \frac{\epsilon}{3}.$$

Then for all x satisfying $|x - x_0| < \delta$, we have

$$\begin{aligned} & |F(x) - F(x_0)| \\ = & |F(x) - F_n(x) + F_n(x) - F_n(x_0) + F_n(x_0) - F(x_0)| \\ \leq & |F(x) - F_n(x)| + |F_n(x) - F_n(x_0)| + |F_n(x_0) - F(x_0)| \end{aligned}$$

$$< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

Thus F is continuous at x_0 . Since x_0 is arbitrary, it follows that F is continuous on I . This finishes the proof of the theorem. \square



EXAMPLE 3.2. Find the pointwise limit F of the sequence

$$F_n(x) = \frac{x^{2n}}{1 + x^{2n}}, \quad x \in [0, 1].$$

Show using Theorem 3.1 that the convergence is not uniform.

SOLUTION. If $0 \leq x < 1$, we have

$$\lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} \frac{x^{2n}}{1 + x^{2n}} = \frac{0}{1 + 0} = 0.$$

If $x = 1$, then $F_n(1) = \frac{1}{2}$. Thus the limiting function $F(x)$ is

$$F(x) = \begin{cases} 0 & 0 \leq x < 1 \\ \frac{1}{2} & x = 1 \end{cases}$$

Because each $F_n(x)$ is continuous but $F(x)$ is not, the sequence of functions $\{F_n(x)\}$ does not converge uniformly on $[0, 1]$ by Theorem 3.1. \square

It is possible that each F_n and $F(x) = \lim_{n \rightarrow \infty} F_n(x)$ are continuous on an interval I , but $\{F_n\}$ does not converge uniformly to $F(x)$.

For the case that I is not closed, an example can be given by $I = [0, 1)$, $F_n(x) = x^n$. (In this case $F(x) = \lim_{n \rightarrow \infty} F_n(x) = 0$ for $x \in I$.)

For the case that I is a closed interval, an example can be given as follows.

EXAMPLE 3.3. Let $F_n(x) = nxe^{-nx^2}$. Show that

- 1) Each $F_n(x)$ is continuous on $I = [0, 1]$.
- 2) $\{F_n(x)\}$ converges pointwise to a continuous function on I .
- 3) $\{F_n(x)\}$ does not converge uniformly on I .

PROOF. Each $F_n(x)$ is continuous because it is a well-defined elementary function on $[0, 1]$.

Let $F(x) = \lim_{n \rightarrow \infty} F_n(x)$ on $[0, 1]$. When $x = 0$, $F_n(0) = 0$ for each n and $F(0) = \lim_{n \rightarrow \infty} 0 = 0$, when $x \neq 0$ (fixed),

$$\begin{aligned} F(x) &= \lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} nxe^{-nx^2} = \lim_{n \rightarrow \infty} \frac{nx}{e^{nx^2}} \\ &= \lim_{n \rightarrow \infty} \frac{x}{e^{nx^2} \cdot x^2} = \lim_{n \rightarrow \infty} \frac{1}{xe^{nx^2}} = 0. \end{aligned}$$

Thus $F(x) = 0$ is continuous on $[0, 1]$.

Now $T_n = \sup_{0 \leq x \leq 1} |F_n(x) - F(x)| = \sup_{0 \leq x \leq 1} F_n(x)$. From

$$\begin{aligned} F'_n(x) &= ne^{-nx^2} + nxe^{-nx^2} \cdot (-2nx) \\ &= ne^{-nx^2}(1 - 2nx^2) = 0, \end{aligned}$$

we have $x = \pm \sqrt{\frac{1}{2n}}$. Since $F_n(0) = 0$ and $F_n(1) = ne^{-n}$,

$$\sup_{0 \leq x \leq 1} F_n(x) = \max_{0 \leq x \leq 1} F_n(x)$$

$$= \max \left\{ 0, \frac{n}{e^n}, n \cdot \frac{1}{\sqrt{2n}} \cdot e^{-n \cdot \frac{1}{2n}} \right\} = \sqrt{\frac{n}{2e}}.$$

Thus $T_n = \sqrt{\frac{n}{2e}}$. Since $\lim_{n \rightarrow \infty} T_n = +\infty$, the sequence $\{F_n\}$ does not converge uniformly to $F(x)$ on $[0, 1]$. \square

COROLLARY 3.4. *Suppose that $\sum_{k=1}^{\infty} f_k$ converges uniformly to a function S on an interval I . Suppose that each f_k is continuous on I . Then S is also continuous on I .*

PROOF. Consider the sequence of partial sums $\{S_n\}$ on I , where we have $S_n = \sum_{k=1}^n f_k$. Then $\{S_n\}$ converges uniformly to S on I . If each f_k is continuous on I , then each S_n is also continuous on I . Then by Theorem 3.1, S is also continuous on I . \square

EXAMPLE 3.5. Is $\sum_{n=1}^{\infty} \frac{x}{n^2 e^{nx}}$, $x \in (0, \infty)$, a continuous function?

SOLUTION. Let $f_n(x) = \frac{x}{n^2 e^{nx}}$.

We find an upper bound of $f_n(x)$. Observe that

$$\begin{aligned} f'_n(x) &= \left(\frac{x}{n^2 e^{nx}} \right)' = \left(\frac{x e^{-nx}}{n^2} \right)' = \frac{e^{-nx} - n x e^{-nx}}{n^2} \\ &= \frac{e^{-nx} \left(\frac{1}{n} - x \right)}{n}. \end{aligned}$$

Thus $f'_n(x) > 0$ for $0 < x < \frac{1}{n}$ and $f'_n(x) < 0$ for $x > \frac{1}{n}$. It follows that $f_n(x)$ is monotone increasing on $(0, \frac{1}{n}]$ and monotone decreasing on $[\frac{1}{n}, +\infty)$. Hence

$$\begin{aligned} \sup_{0 < x < +\infty} f_n(x) &= \max_{0 < x < +\infty} f_n(x) = f_n \left(\frac{1}{n} \right) \\ &= \frac{\frac{1}{n}}{n^2 e^{n \cdot \frac{1}{n}}} = \frac{1}{en^3}. \end{aligned}$$

Let $M_n = \frac{1}{en^3}$. Then $|f_n(x)| \leq M_n$ for $x \in (0, \infty)$.

Since $\sum_{n=1}^{\infty} M_n$ converges by the p -series, the series of

functions $\sum_{n=1}^{\infty} \frac{x}{n^2 e^{nx}}$ converges uniformly by Weierstrass M -test on $(0, \infty)$.

According to Corollary 3.4, the function $\sum_{n=1}^{\infty} \frac{x}{n^2 e^{nx}}$ is continuous on $(0, \infty)$. \square

It is possible that each f_n and $\sum_{n=1}^{\infty} f_n$ are continuous, but $\sum_{n=1}^{\infty} f_n$ does not converge uniformly.

EXAMPLE 3.6. Consider the geometric series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \cdots = \frac{1}{1-x}$$

for $-1 < x < 1$. In this case, each x^n and $\sum_{n=0}^{\infty} x^n$ are continuous on $I = (-1, 1)$. We show that $\sum_{n=0}^{\infty} x^n$ does not converge uniformly by T -test.

Since

$$\begin{aligned} T_n &= \sup_{-1 < x < 1} |x^{n+1} + x^{n+2} + \cdots| \\ &\geq \sup_{0 \leq x < 1} |x^{n+1} + x^{n+2} + \cdots| \\ &= \sup_{0 \leq x < 1} (x^{n+1} + x^{n+2} + \cdots) \geq \sup_{0 \leq x < 1} x^{n+1} = 1, \end{aligned}$$

the sequence $\{T_n\}$ does not tend to 0

and so the series $\sum_{n=0}^{\infty} x^n$ does not converge uniformly on $(-1, 1)$ by the T -test. \square

4. Uniform Convergence and Integration

Before we go on, we first recall some facts about Riemann integrals from MA1102 Calculus I. **Fact 1:**

If $f(x)$ is continuous on a finite closed interval $[a, b]$, then $f(x)$ is **Riemann integrable** on $[a, b]$ (i.e., the


Riemann integral $\int_a^b f(x) dx$ exists in \mathbb{R} .) **Fact 2:**

One always has

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

This follows readily from the following geometric interpretation of the Riemann integral:

$\int_a^b h(x) dx =$ area between the graph of $h(x)$ and the x -axis



See Bartle, Chapter 7 for the above facts and more details on Riemann integrals.

THEOREM 4.1. *Let $\{F_n\}$ be a sequence of continuous functions on a **finite interval** $[a, b]$. Suppose*

that $\{F_n\}$ converges uniformly on $[a, b]$. Then

$$\lim_{n \rightarrow \infty} \int_a^b F_n(x) dx = \int_a^b \left(\lim_{n \rightarrow \infty} F_n(x) \right) dx.$$

PROOF. Let $F(x) = \lim_{n \rightarrow \infty} F_n(x)$. Since each $F_n(x)$ is continuous and $F_n(x)$ converges uniformly on $[a, b]$, the function $F(x)$ is continuous on $[a, b]$ and the integral $\int_a^b F(x) dx$ exists.

By definition, given any $\epsilon > 0$, there exists N such that

$$|F_n(x) - F(x)| < \frac{\epsilon}{2(b-a)}$$

for all $n > N$ and $x \in [a, b]$. Thus

$$F_n(x) - \frac{\epsilon}{2(b-a)} < F(x) < F_n(x) + \frac{\epsilon}{2(b-a)}$$

for all $x \in [a, b]$ and $n > N$.

It follows that, for $n > N$,

$$\begin{aligned} \int_a^b \left(F_n(x) - \frac{\epsilon}{2(b-a)} \right) dx &\leq \int_a^b F(x) dx \\ &\leq \int_a^b \left(F_n(x) + \frac{\epsilon}{2(b-a)} \right) dx \end{aligned}$$

and so

$$\int_a^b F_n(x) dx - \frac{\epsilon}{2} \leq \int_a^b F(x) dx \leq \int_a^b F_n(x) dx + \frac{\epsilon}{2}$$

or

$$\left| \int_a^b F_n(x) dx - \int_a^b F(x) dx \right| \leq \frac{\epsilon}{2} < \epsilon.$$

Thus

$$\lim_{n \rightarrow \infty} \int_a^b F_n(x) dx = \int_a^b F(x) dx = \int_a^b \lim_{n \rightarrow \infty} F_n(x) dx,$$

which is the assertion.

□

EXAMPLE 4.2. Compute, justifying your answer,

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{\sin nx}{n + x^2} dx.$$

SOLUTION. Let $F_n(x) = \frac{\sin nx}{n + x^2}$. Then the limiting function

$$F(x) = \lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} \frac{\sin nx}{n + x^2} = 0$$

for any given $0 \leq x \leq 1$, by the Squeeze theorem, because

$$-\frac{1}{n} \leq \frac{\sin nx}{n + x^2} \leq \frac{1}{n}$$

and $\lim_{n \rightarrow \infty} \frac{1}{n} = -\lim_{n \rightarrow \infty} \frac{1}{n} = 0$. Since

$$T_n = \sup_{0 \leq x \leq 1} |F_n(x) - F(x)| = \sup_{0 \leq x \leq 1} \left| \frac{\sin nx}{n + x^2} \right| \leq \frac{1}{n},$$

$\lim_{n \rightarrow \infty} T_n = 0$ by the Squeeze Theorem and so the sequence of functions $\{F_n\}$ converges uniformly to $F(x)$.

Thus

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{\sin nx}{n + x^2} dx = \int_0^1 \lim_{n \rightarrow \infty} \frac{\sin nx}{n + x^2} dx = \int_0^1 0 dx = 0.$$

□

COROLLARY 4.3. *Suppose that $\sum_{k=1}^{\infty} f_k$ converges uniformly on a finite interval $[a, b]$ such that each f_k is continuous on $[a, b]$. Then*

$$\int_a^b \sum_{k=1}^{\infty} f_k(x) dx = \sum_{k=1}^{\infty} \int_a^b f_k(x) dx.$$

PROOF. By definition, the sequence of partial sums $S_n = \sum_{k=1}^n f_k$ converges uniformly on $[a, b]$. Since each f_k is continuous on $[a, b]$, then each partial sum S_n is also continuous on $[a, b]$. Thus

$$\begin{aligned} \int_a^b \sum_{k=1}^{\infty} f_k dx &= \int_a^b \lim_{n \rightarrow \infty} S_n(x) dx \\ &= \lim_{n \rightarrow \infty} \int_a^b S_n(x) dx = \lim_{n \rightarrow \infty} \int_a^b \sum_{k=1}^n f_k(x) dx \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \int_a^b f_k(x) dx = \sum_{k=1}^{\infty} \int_a^b f_k(x) dx. \end{aligned}$$

□

By using this theorem, we have the following amazing formula.

EXAMPLE 4.4. *Show that*

$$\ln 2 = \sum_{n=1}^{\infty} \frac{1}{n \cdot 2^n} = \frac{1}{2} + \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} + \cdots$$

PROOF. Since $|x^{n-1}| \leq \left(\frac{1}{2}\right)^{n-1}$ for $0 \leq x \leq \frac{1}{2}$ and the geometric series $\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n-1}$ converges, the series

$\sum_{n=1}^{\infty} x^{n-1}$ converges uniformly to $\frac{1}{1-x}$ on $[0, \frac{1}{2}]$ by the Weierstrass M -test. Thus we have

$$\begin{aligned} \int_0^{\frac{1}{2}} \frac{1}{1-x} dx &= \int_0^{\frac{1}{2}} \sum_{n=1}^{\infty} x^{n-1} dx \\ &= \sum_{n=1}^{\infty} \int_0^{\frac{1}{2}} x^{n-1} dx = \sum_{n=1}^{\infty} \frac{x^n}{n} \Big|_0^{\frac{1}{2}} = \sum_{n=1}^{\infty} \frac{1}{n \cdot 2^n}. \end{aligned}$$

Since

$$\begin{aligned} \int_0^{\frac{1}{2}} \frac{1}{1-x} dx &= -\ln(1-x) \Big|_0^{\frac{1}{2}} = -\ln\left(1 - \frac{1}{2}\right) \\ &= -\ln\left(\frac{1}{2}\right) = \ln 2, \end{aligned}$$

we obtain the formula

$$\ln 2 = \sum_{n=1}^{\infty} \frac{1}{n \cdot 2^n} = \frac{1}{2} + \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} + \cdots .$$

□

REMARK 4.5. Example 4.4 gives a way to estimate the number $\ln 2$ because the remainder

$$\begin{aligned} R_n &= \sum_{k=n+1}^{\infty} \frac{1}{k \cdot 2^k} = \frac{1}{(n+1)2^{n+1}} + \frac{1}{(n+2)2^{n+2}} + \cdots \\ &< \frac{1}{(n+1)2^{n+1}} + \frac{1}{(n+1)2^{n+2}} + \frac{1}{(n+1)2^{n+3}} + \cdots \\ &= \frac{1}{(n+1)2^{n+1}} \left(1 + \frac{1}{2} + \frac{1}{2^2} + \cdots \right) \\ &= \frac{1}{(n+1)2^{n+1}} \cdot \frac{1}{1 - \frac{1}{2}} = \frac{1}{(n+1)2^n}. \end{aligned}$$

For instance,

$$\ln 2 \approx \frac{1}{2} + \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} + \cdots + \frac{1}{10 \cdot 2^{10}}$$

with error less than $\frac{1}{11 \cdot 2^{10}} = \frac{1}{11264}$.

5. Uniform Convergence and Differentiation

THEOREM 5.1. *Let $\{F_n\}$ be a sequence of functions on $[a, b]$ such that*

- (i) *each F'_n exists and is continuous on $[a, b]$,*
- (ii) *$\{F_n\}$ converges pointwise to a function F on $[a, b]$, and*
- (iii) *$\{F'_n\}$ converges uniformly on $[a, b]$.*

Then F is differentiable on $[a, b]$, and for all $x \in [a, b]$,

$$F'(x) = \lim_{n \rightarrow \infty} F'_n(x),$$

i.e.
$$\frac{d}{dx} \left(\lim_{n \rightarrow \infty} F_n(x) \right) = \lim_{n \rightarrow \infty} \left(\frac{d}{dx} F_n(x) \right).$$

Remark. Here the differentiability and continuity at the endpoints a and b refer to the one sided derivatives and limits respectively.

PROOF. By (iii), $\{F'_n\}$ converges uniformly to a function, denoted by g , on $[a, b]$, where

$$g(x) = \lim_{n \rightarrow \infty} F'_n(x)$$

for all $x \in [a, b]$.

By (i), since each F'_n is continuous on $[a, b]$, F'_n is also Riemann integrable on $[a, b]$, and by the fundamental theorem of calculus,

$$\int_a^x F'_n(t) dt = F_n(x) - F_n(a) \quad \text{for all } x \in [a, b].$$

Letting $n \rightarrow \infty$, for all $x \in [a, b]$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_a^x F'_n(t) dt &= \lim_{n \rightarrow \infty} (F_n(x) - F_n(a)) = F(x) - F(a) \\ \underline{\underline{F'_n \text{ uniform convergence on } [a, x] \subseteq [a, b]}} &\int_a^x \lim_{n \rightarrow \infty} F'_n(t) dt = \int_a^x g(t) dt. \end{aligned}$$

Thus

$$F(x) = F(a) + \int_a^x g(t) dt$$

and so

$$F'(x) = \frac{d}{dx} \int_a^x g(t) dt = g(x)$$

by the fundamental theorem of calculus. In other words

$$\frac{d}{dx} \left(\lim_{n \rightarrow \infty} F_n(x) \right) = \lim_{n \rightarrow \infty} \left(\frac{d}{dx} F_n(x) \right)$$

and hence the result. \square

REMARK 5.2. *By inspecting the proof, Theorem 5.1 still holds when the closed interval $[a, b]$ is replaced by (a, b) , $(a, b]$ or $[a, b)$.*

COROLLARY 5.3. *Let $\sum_{k=1}^{\infty} f_k$ be a series of functions on $[a, b]$ such that*

- (i) *each f'_k exists and continuous on $[a, b]$,*
- (ii) $\sum_{k=1}^{\infty} f_k$ *converges pointwise on $[a, b]$, and*
- (iii) $\sum_{k=1}^{\infty} f'_k$ *converges **uniformly** on $[a, b]$.*

Then

$$\frac{d}{dx} \left(\sum_{k=1}^{\infty} f_k(x) \right) = \sum_{k=1}^{\infty} \frac{d}{dx} f_k(x).$$

PROOF.

$$\begin{aligned} \frac{d}{dx} \left(\sum_{k=1}^{\infty} f_k(x) \right) &= \frac{d}{dx} \lim_{n \rightarrow \infty} \sum_{k=1}^n f_k(x) \\ &\stackrel{\text{check conditions (i)-(iii)}}{=} \lim_{n \rightarrow \infty} \frac{d}{dx} \left(\sum_{k=1}^n f_k(x) \right) \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n f'_k(x) = \sum_{k=1}^{\infty} \frac{d}{dx} f_k(x). \end{aligned}$$

□

Example. Let $S(x) = \sum_{k=1}^{\infty} \frac{\sin kx}{2^k}$. Let $f_k(x) = \frac{\sin kx}{2^k}$.

Then

(i) each f'_k exists and continuous on \mathbb{R} .

(ii) $\sum_{k=1}^{\infty} f_k$ converges pointwise on \mathbb{R} because

$$|f_k(x)| \leq \frac{1}{2^k}$$

and $\sum_{k=1}^{\infty} \frac{1}{2^k}$ converges.

(iii)

$$\sum_{k=1}^{\infty} f'_k = \sum_{k=1}^{\infty} \frac{k \cos kx}{2^k}$$

converges **uniformly** on \mathbb{R} by the Weierstrass M -test because

$$\left| \frac{k \cos kx}{2^k} \right| \leq \frac{k}{2^k}$$

and $\sum_{k=1}^{\infty} \frac{k}{2^k}$ converges.

Thus

$$S'(x) = \sum_{k=1}^{\infty} \frac{k \cos kx}{2^k}.$$

6. Power Series

6.1. Power Series.

DEFINITION 6.1. A **power series** in x is of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \cdots$$

EXAMPLE 6.2. Below are some examples

1.
$$\sum_{n=0}^{\infty} (n+1)x^n = 1 + 2x + 3x^2 + 4x^3 + 5x^4 + \cdots$$
2.
$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

DEFINITION 6.3. A **power series** in $x - x_0$ is of the form

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n = a_0 + a_1(x - x_0) + a_2(x - x_0)^2 + \cdots$$

EXAMPLE 6.4. Here are some examples.

$$1. \quad \sum_{n=0}^{\infty} (x - 1)^n = 1 + (x - 1) + (x - 1)^2 + \cdots .$$

$$2. \quad \sum_{n=1}^{\infty} n^2(x + 2)^n = (x + 2) + 2^2(x + 2)^2 + 3^2(x + 2)^3 + \cdots .$$

Warning. Don't expand out the terms $a_n(x - x_0)^n$

in the power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ because, when you

rearrange terms in an (infinite) series, you may get different values. (For partial sums, you can expand out, if it is necessary, because there are only finitely many terms.)

Question: Given a power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$,

when does it converge and when does it diverge? In

other words, what is the domain of the function $\sum_{n=0}^{\infty} a_n(x -$

$x_0)^n$. We are going to answer this question.

6.2. Radius of Convergence.

DEFINITION 6.5. Given a power series $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, the radius of convergence R is defined by

$$R = \frac{1}{\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|}}.$$

If $\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \infty$, we take $R = 0$, and if $\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} = 0$, we set $R = \infty$. If $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}$ exists, R is also given by

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}}$$

Remark. Recall that

$$\underline{\lim}_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} \leq \underline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} \leq \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} \leq \overline{\lim}_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}.$$

If $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}$ exists, then $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}$ exists with

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}.$$

EXAMPLE 6.6. *What is the radius of convergence for the series*

$$1 + \frac{x}{3} + \frac{x^2}{4^2} + \frac{x^3}{3^3} + \frac{x^4}{4^4} + \frac{x^5}{3^5} + \frac{x^6}{4^6} + \dots$$

SOLUTION. Since

$$a_n = \begin{cases} \frac{1}{4^{2k}} & n = 2k \\ \frac{1}{3^{2k-1}} & n = 2k - 1, \end{cases}$$

we have

$$\sqrt[n]{|a_n|} = \begin{cases} \frac{1}{4} & n = 2k \\ \frac{1}{3} & n = 2k - 1. \end{cases}$$

Thus $\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \frac{1}{3}$ and so the radius of convergence

$$R = \frac{1}{\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|}} = \frac{1}{\frac{1}{3}} = 3.$$

□

EXAMPLE 6.7. Find the radius of convergence of the power series $\sum_{n=0}^{\infty} \frac{(4x+3)^n}{n^3}$

SOLUTION. Observe that

$$\sum_{n=1}^{\infty} \frac{(4x+3)^n}{n^3} = \sum_{n=1}^{\infty} \frac{4^n}{n^3} \cdot \left(x + \frac{3}{4}\right)^n.$$

Thus

$$\begin{aligned} R &= \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{4^{n+1} \cdot n^3}{(n+1)^3 \cdot 4^n}} \\ &= \frac{1}{\lim_{n \rightarrow \infty} \frac{4}{\left(1 + \frac{1}{n}\right)^3}} = \frac{1}{4}. \end{aligned}$$

□

THEOREM 6.8. *Given any power series $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ with **radius of convergence** R , $0 \leq R \leq \infty$, then the series $\sum_{k=0}^{\infty} a_k(x - x_0)^k$*

- (i) **converges absolutely** for all x with $|x - x_0| < R$, and
- (ii) **diverges** for all x with $|x - x_0| > R$.

Remark: At the **(two) end points** $|x - x_0| = R$, that is, $x = x_0 \pm R$, the power series **possibly converges** and **possibly diverges**.

PROOF. By definition, the radius of convergence

$$R = \frac{1}{\overline{\lim}_{k \rightarrow \infty} |a_k|^{\frac{1}{k}}}.$$

Assertion (i) follows from the root test because, from

$$\begin{aligned} \overline{\lim}_{k \rightarrow \infty} |a_k(x - x_0)^k|^{\frac{1}{k}} &= \overline{\lim}_{k \rightarrow \infty} |a_k|^{\frac{1}{k}} \cdot |x - x_0| \\ &= |x - x_0| \cdot \overline{\lim}_{k \rightarrow \infty} |a_k|^{\frac{1}{k}} \\ &= |x - x_0| \cdot \frac{1}{R} < R \cdot \frac{1}{R} < 1, \end{aligned}$$

the series $\sum_{k=0}^{\infty} |a_k(x - x_0)^k|$ converges.

Next we are going to prove (ii) by contradiction.

Suppose that $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ converges at a point x with $|x - x_0| > R$.

Then by Theorem 1.7, we have

$$\lim_{k \rightarrow \infty} a_k(x - x_0)^k = 0.$$

Let $\epsilon = 1$. Then there exists N such that

$$\begin{aligned} & |a_k(x - x_0)^k - 0| < 1 \quad \text{for all } k > N \\ \implies & |a_k(x - x_0)^k|^{\frac{1}{k}} < 1 \quad \text{for all } k > N \\ \implies & |a_k|^{\frac{1}{k}} < \frac{1}{|x - x_0|} \quad \text{for all } k > N \\ \implies & \sup_{n \geq k} |a_n|^{\frac{1}{n}} \leq \frac{1}{|x - x_0|} \quad \text{for all } n > N \\ \implies & \overline{\lim}_{k \rightarrow \infty} |a_k|^{\frac{1}{k}} \leq \frac{1}{|x - x_0|} \\ \implies & \frac{1}{R} \leq \frac{1}{|x - x_0|} < \frac{1}{R}, \end{aligned}$$

which is a contradiction.

Hence we must have $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ diverges at each x satisfying $|x - x_0| > R$. □

6.3. Interval of convergence. In view of Theorem 6.8, for a power series $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ with radius of convergence R , the set of points at which $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ is convergent form an interval called the **interval of convergence**, which must be either

$$(x_0 - R, x_0 + R), \quad (x_0 - R, x_0 + R], \\ [x_0 - R, x_0 + R) \quad \text{or} \quad [x_0 - R, x_0 + R].$$

EXAMPLE 6.9. Find the interval of convergence of the power series.

$$(i) \sum_{n=1}^{\infty} \frac{(x-2)^n}{n^2} \quad (ii) \sum_{n=1}^{\infty} \frac{(x-2)^n}{n} \quad (iii) \sum_{n=1}^{\infty} n(x-2)^n$$

SOLUTION. (i). First we find the radius of convergence

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{n^2}{(n+1)^2}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{1}{(1 + \frac{1}{n})^2}} = 1.$$

Next we check the ending-points $x_0 \pm R = 2 \pm 1 = 1, 3$. When $x = 1$, the series is $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$, which is convergent by Example 3.3. When $x = 3$, the series is

$\sum_{n=1}^{\infty} \frac{1}{n^2}$, which is convergent by the p -series. Thus the interval of convergence is $[1, 3]$.

$$(ii). \sum_{n=1}^{\infty} \frac{(x-2)^n}{n}.$$

The radius of convergence is

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{n}{n+1}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}}} = 1.$$

Now we check the ending-points $x_0 \pm R = 1, 3$.

When $x = 1$, the series is $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$, which is convergent by Example 3.3. When $x = 3$, the series is $\sum_{n=1}^{\infty} \frac{1}{n}$, which is divergent by the p -series. Thus the interval of convergence is $[1, 3)$.

$$(iii). \sum_{n=1}^{\infty} n(x-2)^n.$$

The radius of convergence is

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}} = \frac{1}{\lim_{n \rightarrow \infty} \frac{n+1}{n}} = \frac{1}{\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)} = 1.$$

Now we check the ending-points $x_0 \pm R = 1, 3$.

When $x = 1$, the series is $\sum_{n=1}^{\infty} n(-1)^n$, and when $x = 3$,

the series is $\sum_{n=1}^{\infty} n$. Both of these series are divergent by the divergence test. Thus the interval of convergence is $(1, 3)$. \square

Remarks on radius and interval of convergence of power series:

Keep in mind that **interval of convergence** is just the **domain of the power series**, and **radius of convergence** is just distance from the center to the ending points. (So it is just the **half-length of the interval**.)

For some power series, that is hard to be written in the standard form, we can simply try to find the domain for getting interval or radius of convergence.

Example. Find the radius and interval of the power

series
$$\sum_{n=0}^{\infty} \frac{(x-1)^{n^2}}{n}.$$

SOLUTION. By applying the root test for general series, let

$$\begin{aligned}
 l &= \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\left| \frac{(x-1)^{n^2}}{n} \right|} = \lim_{n \rightarrow \infty} \frac{|x-1|^n}{\sqrt[n]{n}} \\
 &= \begin{cases} +\infty & |x-1| > 1 \\ 1 & |x-1| = 1 \\ 0 & |x-1| < 1. \end{cases}
 \end{aligned}$$

Thus the series converges (absolutely) when $|x-1| < 1$, and diverges when $|x-1| > 1$. Check the ending points $|x-1| = 1$, that is, $x = 0, 2$. When $x = 0$,

the series is $\sum_{n=0}^{\infty} \frac{(-1)^{n^2}}{n} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n}$, (where $(-1)^{n^2} =$

$(-1)^n$ by checking n odd or even), and it converges by the alternating series test. When $x = 2$, the series is

$\sum_{n=0}^{\infty} \frac{1}{n}$ which is divergent by the p -series.

Hence the interval of convergence is $[0, 2)$ and the radius of convergence is 1.

□

6.4. Uniform Convergence of Power Series.

THEOREM 6.10 (Uniform Convergence Theorem).
Suppose that a power series

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n$$

converges when $x = c, d$. Then the series of functions $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges uniformly on the closed interval $[c, d]$.

PROOF. Omitted. See the packed lecture notes. \square

Remark. The theorem says that if a **power series** converges at the ending points of a closed interval $[c, d]$, then it converges uniformly on $[c, d]$. This is a **special property** of power series on uniform convergence.

COROLLARY 6.11. Let $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ be a power series such that it is **convergent** when $x = c, d$. Then

$$\int_c^d \sum_{n=0}^{\infty} a_n(x - x_0)^n dx = \sum_{n=0}^{\infty} \int_c^d a_n(x - x_0)^n dx.$$

PROOF. Since $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ converges when $x = c, d$, it converges uniformly on $[c, d]$ and so

$$\int_c^d \sum_{n=0}^{\infty} a_n(x - x_0)^n dx = \sum_{n=0}^{\infty} \int_c^d a_n(x - x_0)^n dx.$$

□

COROLLARY 6.12 (Abel Theorem). *Let $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ be a power series of radius of convergence $R > 0$.*

(a). *If $\sum_{n=0}^{\infty} a_n R^n$ **converges**, (i.e. the power series converges when $x = x_0 + R$), then*

$$\begin{aligned} \lim_{x \rightarrow (x_0 + R)^-} \sum_{n=0}^{\infty} a_n (x - x_0)^n &= \sum_{n=0}^{\infty} \lim_{x \rightarrow (x_0 + R)^-} a_n (x - x_0)^n \\ &= \sum_{n=0}^{\infty} a_n R^n. \end{aligned}$$

(b). *If $\sum_{n=0}^{\infty} a_n (-R)^n$ **converges**, (i.e. the power series converges when $x = x_0 - R$), then*

$$\begin{aligned} \lim_{x \rightarrow (x_0 - R)^+} \sum_{n=0}^{\infty} a_n (x - x_0)^n &= \sum_{n=0}^{\infty} \lim_{x \rightarrow (x_0 - R)^+} a_n (x - x_0)^n \\ &= \sum_{n=0}^{\infty} a_n (-R)^n. \end{aligned}$$

PROOF. (a). Since the power series

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n$$

converges when $x = x_0, x_0 + R$, it converges uniformly on $[x_0, x_0 + R]$ and so

$$\begin{aligned} \lim_{x \rightarrow (x_0+R)^-} \sum_{n=0}^{\infty} a_n(x-x_0)^n &= \sum_{n=0}^{\infty} \lim_{x \rightarrow (x_0+R)^-} a_n(x-x_0)^n \\ &= \sum_{n=0}^{\infty} a_n R^n. \end{aligned}$$

The proof of (b) is similar to that of (a). □

EXAMPLE 6.13. From the geometric series $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$ for $|x| < 1$, we have

$$\frac{1}{1+t} = \sum_{n=0}^{\infty} (-1)^n t^n$$

by letting $x = -t$. For any $x \in (-1, 1)$, we have

$$\begin{aligned} \ln(1+x) &= \int_0^x \frac{1}{1+t} dt \\ &= \sum_{n=0}^{\infty} (-1)^n \int_0^x t^n dt = \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1}. \end{aligned}$$

Since $\sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1}$ converges when $x = 1$, we have

$$\begin{aligned} \sum_{n=0}^{\infty} (-1)^n \frac{1}{n+1} &= \lim_{x \rightarrow 1^-} \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1} \\ &= \lim_{x \rightarrow 1^-} \ln(1+x) = \ln 2. \end{aligned}$$

In other words,

$$\ln 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots .$$

EXAMPLE 6.14. From the geometric series $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$ for $|x| < 1$, we have

$$\frac{1}{1+t^2} = \sum_{n=0}^{\infty} (-1)^n t^{2n}$$

by letting $x = -t^2$. For any $x \in (-1, 1)$, we have

$$\begin{aligned} \arctan x &= \int_0^x \frac{1}{1+t^2} dt \\ &= \sum_{n=0}^{\infty} (-1)^n \int_0^x t^{2n} dt = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}. \end{aligned}$$

Since $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$ converges when $x = \pm 1$, it converges uniformly on $[-1, 1]$ and so

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \text{for all } |x| \leq 1.$$

In particular,

$$\frac{\pi}{4} = \arctan 1 = \sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots.$$

EXAMPLE 6.15. From

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \text{for all } |x| \leq 1,$$

we have

$$\arctan x^2 = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2(2n+1)}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1}$$

for $|x| \leq 1$ and so

$$\begin{aligned} \int_0^1 \arctan x^2 dx &= \sum_{n=0}^{\infty} (-1)^n \int_0^1 \frac{x^{4n+2}}{2n+1} dx \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)(4n+3)}. \end{aligned}$$

7. Differentiation of Power Series

LEMMA 7.1. *Let $\{a_n\}$ and $\{b_n\}$ be sequences such that $a_n \geq 0$, $\lim_{n \rightarrow \infty} b_n$ exists with $\lim_{n \rightarrow \infty} b_n \neq 0$. Then*

$$\overline{\lim}_{n \rightarrow \infty} a_n b_n = \overline{\lim}_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n.$$

PROOF. Omitted □

THEOREM 7.2. *Suppose that $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ has radius of convergence $R > 0$, and*

$$f(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n, \quad |x - x_0| < R.$$

Then

(a). *The power series $\sum_{n=1}^{\infty} n a_n(x - x_0)^{n-1}$ has radius of convergence R , and*

(b). $f'(x) = \sum_{n=1}^{\infty} n a_n(x - x_0)^{n-1}$ for $|x - x_0| < R$.

PROOF. (a). By Lemma 7.1,

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} |na_n|^{\frac{1}{n}} &= \overline{\lim}_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} \cdot n^{\frac{1}{n}} = \overline{\lim}_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} \cdot \lim_{n \rightarrow \infty} \sqrt[n]{n} \\ &= \overline{\lim}_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} \cdot 1 = \overline{\lim}_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} = \frac{1}{R}.\end{aligned}$$

Thus the power series

$$\begin{aligned}\sum_{n=0}^{\infty} na_n(x - x_0)^n &= \sum_{n=1}^{\infty} na_n(x - x_0)^n \\ &= (x - x_0) \cdot \sum_{n=1}^{\infty} na_n(x - x_0)^{n-1}\end{aligned}$$

has radius of convergence R and so has $\sum_{n=1}^{\infty} na_n(x - x_0)^{n-1}$.

(b). For any ρ with $0 < \rho < R$, the series of functions

$$\sum_{n=1}^{\infty} na_n(x - x_0)^{n-1}$$

converges uniformly on $|x - x_0| \leq \rho$ by the Uniform Convergence Theorem because the closed interval $[x_0 - \rho, x_0 + \rho] \subseteq (x_0 - R, x_0 + R)$. Thus we can do derivatives term-by-term and hence the result. \square

Remark. The formula $f'(x) = \sum_{n=1}^{\infty} na_n(x - x_0)^{n-1}$ need not hold at the end points $x = x_0 \pm R$ in general even if the interval of convergence of $\sum_{n=0}^{\infty} a_n(x - x_0)^n$ is $[x_0 - R, x_0 + R]$.

EXAMPLE 7.3. From Example 6.14,

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \text{for all } |x| \leq 1.$$

But

$$\frac{1}{1+x^2} = (\arctan x)' = \sum_{n=0}^{\infty} (-1)^n x^{2n}$$

only holds for $|x| < 1$ because when $x = \pm 1$, the right hand side diverges (and the left hand side $= \frac{1}{2}$).

COROLLARY 7.4. *Suppose that $\sum_{k=0}^{\infty} a_k(x-x_0)^k$ has radius of convergence $R > 0$. Let*

$$f(x) = \sum_{k=0}^{\infty} a_k(x-x_0)^k$$

on $|x-x_0| < R$.

Then $f(x)$ has **derivatives of all orders** on $|x-x_0| < R$, and for each n ,

$$(6) \quad f^{(n)}(x) = \sum_{k=n}^{\infty} k(k-1)(k-2) \cdots (k-n+1) a_k (x-x_0)^{k-n}.$$

In particular,

$$(7) \quad a_k = \frac{f^{(k)}(x_0)}{k!} \quad \text{for all } k.$$

(i.e. we have $f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k$.)

PROOF. The result is obtained by successively applying the previous theorem to f , f' , f'' , and etc. Equation 7 follows by setting $x = x_0$ in Equation 6, that is,

$$\begin{aligned} f^{(n)}(x) &= n!a_n + (n+1)n \cdots 2a_{n+1}(x-x_0) \\ &+ (n+2)(n+1) \cdots 3a_{n+2}(x-x_0)^2 + \cdots . \end{aligned}$$

□

EXAMPLE 7.5. From the geometric series

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \cdots \quad |x| < 1,$$

we have

$$\frac{1}{(1-x)^2} = \left(\frac{1}{1-x} \right)' = \sum_{n=1}^{\infty} nx^{n-1} = 1 + 2x + 3x^2 + 4x^3 + \cdots$$

for $|x| < 1$, and so

$$\frac{x}{(1-x)^2} = \sum_{n=1}^{\infty} nx^n = x + 2x^2 + 3x^3 + 4x^4 + \cdots \quad |x| < 1.$$

By letting $x = \frac{1}{2}$, we have

$$\sum_{n=1}^{\infty} \frac{n}{2^n} = \frac{\frac{1}{2}}{\left(1 - \frac{1}{2}\right)^2} = 2$$

As an application, we give a proof of binomial series. Let a be any real number. The binomial number, a chooses n , is defined by

$$\binom{a}{n} = \frac{a(a-1)(a-2)\cdots(a-n+1)}{n!}$$

for positive integers n . For instance,

$$\begin{aligned} \binom{\frac{1}{2}}{1} &= \frac{1}{2}, & \binom{\frac{1}{2}}{2} &= \frac{\frac{1}{2} \cdot (\frac{1}{2} - 1)}{2!} = -\frac{1}{8}, \\ \binom{\frac{1}{2}}{3} &= \frac{\frac{1}{2} \cdot (\frac{1}{2} - 1) (\frac{1}{2} - 2)}{3!} = \frac{\frac{1}{2} \cdot (-\frac{1}{2}) \cdot (-\frac{3}{2})}{6} = \frac{1}{16}. \end{aligned}$$

We also use the convention that $\binom{a}{0} = 1$ for any a .

THEOREM 7.6 (Binomial Series). *Let a be any real constant. Then*

$$\begin{aligned} (1+x)^a &= 1 + ax + \frac{a(a-1)}{2!}x^2 + \cdots \\ &= 1 + \sum_{n=1}^{\infty} \binom{a}{n} x^n = \sum_{n=0}^{\infty} \binom{a}{n} x^n \end{aligned}$$

for $|x| < 1$.

PROOF. First we are going to do derivatives term-by-term for the power series. Let

$$f(x) = 1 + \sum_{n=1}^{\infty} \binom{a}{n} x^n.$$

Since

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{\left| \binom{a}{n+1} \right|}{\left| \binom{a}{n} \right|} \\ &= \lim_{n \rightarrow \infty} \frac{|a| \cdot |a-1| \cdots |a-n|}{(n+1)!} \\ &= \lim_{n \rightarrow \infty} \frac{|a| \cdot |a-1| \cdots |a-n+1|}{n!} \\ &= \lim_{n \rightarrow \infty} \frac{|a| \cdot |a-1| \cdots |a-n| \cdot n!}{(n+1)! \cdot |a| \cdot |a-1| \cdots |a-n+1|} \\ &= \lim_{n \rightarrow \infty} \frac{|a-n|}{n+1} = 1, \end{aligned}$$

the power series $f(x) = 1 + \sum_{n=1}^{\infty} \binom{a}{n} x^n$ converges on $(-1, 1)$ and so

$$f'(x) = \sum_{n=1}^{\infty} \binom{a}{n} n x^{n-1}$$

for $-1 < x < 1$.

Next we are going to set up a differential equation that

$$(1+x)f'(x) = af(x).$$

(Note. Since the goal is to show that $f(x) = (1+x)^a$, this equation is observed from that, if $y = (1+x)^a$, then $y' = a(1+x)^{a-1}$ and so $(1+x)y' = (1+x)^a = y$.)

$$\begin{aligned}
 f'(x) &= \sum_{n=1}^{\infty} \binom{a}{n} nx^{n-1} \\
 &= \sum_{n=1}^{\infty} \frac{a(a-1)\cdots(a-n+1) \cdot n}{n!} x^{n-1} \\
 &= \sum_{n=1}^{\infty} \frac{a(a-1)\cdots(a-n+1)}{(n-1)!} x^{n-1} \\
 &= a \sum_{n=1}^{\infty} \binom{a-1}{n-1} x^{n-1} \quad \text{and}
 \end{aligned}$$

$$\begin{aligned}
 (1+x)f'(x) &= f'(x) + xf'(x) \\
 &= a \sum_{n=1}^{\infty} \binom{a-1}{n-1} x^{n-1} + a \sum_{n=1}^{\infty} \binom{a-1}{n-1} x^n \\
 &= a \sum_{n=0}^{\infty} \binom{a-1}{n} x^n + a \sum_{n=1}^{\infty} \binom{a-1}{n-1} x^n \\
 &= a \left\{ 1 + \sum_{n=1}^{\infty} \left[\binom{a-1}{n} + \binom{a-1}{n-1} \right] x^n \right\} \\
 &= a \left[1 + \sum_{n=1}^{\infty} \binom{a}{n} x^n \right] = af(x),
 \end{aligned}$$

where

$$\begin{aligned}
 & \binom{a-1}{n} + \binom{a-1}{n-1} \\
 &= \frac{(a-1)(a-2)\cdots(a-1-n+1)}{n!} \\
 & \quad + \frac{(a-1)(a-2)\cdots(a-1-n+2)}{(n-1)!} \\
 &= \frac{(a-1)(a-2)\cdots(a-1-n+2)}{n!} (a-1-n+1+n) \\
 &= \frac{a(a-1)\cdots(a-n+1)}{n!} = \binom{a}{n}.
 \end{aligned}$$

Our last step is to solve the differential equation

$$(1+x)f'(x) = af(x).$$

Let $y = f(x)$. Then we obtain the differential equation

$$\begin{aligned}
 (1+x)\frac{dy}{dx} &= ay & \frac{dy}{y} &= \frac{a dx}{1+x} \\
 \implies \int \frac{dy}{y} &= \int \frac{a dx}{1+x} \\
 \implies \ln|y| &= a \ln|1+x| + A = \ln|1+x|^a + A \\
 \implies |y| &= e^{\ln|y|} = e^A |1+x|^a \\
 \implies y &= C|1+x|^a,
 \end{aligned}$$

where $C = \pm e^A$ is a constant. By putting $x = 0$,

$$C = (1+0)^a = y(0) = 1 + \sum_{n=1}^{\infty} \binom{a}{n} 0^n = 1.$$

Thus $y = |1 + x|^a$ or

$$(1 + x)^a = 1 + \sum_{n=1}^{\infty} \binom{a}{n} x^n$$

for $|x| < 1$ because $1 + x > 0$ when $|x| < 1$. □

For instance,

$$\begin{aligned}\sqrt{1+x} &= (1+x)^{\frac{1}{2}} = \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} x^k \\ &= 1 + \frac{1}{2}x + \frac{\binom{\frac{1}{2}}{2} \left(-\frac{1}{2}\right)}{2!} x^2 + \dots \\ \sqrt{1-x^3} &= (1-x^3)^{\frac{1}{2}} = \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} (-x^3)^k \\ &= 1 - \frac{1}{2}x^3 + \frac{\binom{\frac{1}{2}}{2} \left(-\frac{1}{2}\right)}{2!} x^6 + \dots \\ \sqrt{4.1} &= \left(4 + \frac{1}{10}\right)^{\frac{1}{2}} = 2 \left(1 + \frac{1}{40}\right)^{\frac{1}{2}} \\ &= 2 \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} \left(\frac{1}{40}\right)^k.\end{aligned}$$

EXAMPLE 7.7. Evaluate $\sqrt{4.1}$ with error less than 0.001.

SOLUTION.

$$\begin{aligned}\sqrt{4.1} &= \left(4 + \frac{1}{10}\right)^{\frac{1}{2}} = 2 \left(1 + \frac{1}{40}\right)^{\frac{1}{2}} = 2 \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} \left(\frac{1}{40}\right)^k \\ &= 2 + 2 \sum_{k=1}^{\infty} \binom{\frac{1}{2}}{k} \left(\frac{1}{40}\right)^k.\end{aligned}$$

Now

$$\begin{aligned} \binom{\frac{1}{2}}{k} &= \frac{\frac{1}{2} \cdot \left(\frac{1}{2} - 1\right) \cdots \left(\frac{1}{2} - k + 1\right)}{k!} \\ &= (-1)^{k+1} \frac{\frac{1}{2} \cdot \left(1 - \frac{1}{2}\right) \cdots \left(k - \frac{1}{2} - 1\right)}{k!} \end{aligned}$$

for $k \geq 2$. Let

$$b_k = \frac{\frac{1}{2} \cdot \left(1 - \frac{1}{2}\right) \cdots \left(k - \frac{1}{2} - 1\right)}{k!} \left(\frac{1}{40}\right)^k.$$

Then, for $k \geq 2$, we have $b_k \geq 0$,

$$\begin{aligned} \frac{b_{k+1}}{b_k} &= \frac{\frac{\frac{1}{2} \cdot \left(1 - \frac{1}{2}\right) \cdots \left(k - \frac{1}{2} - 1\right) \cdot \left(k - \frac{1}{2}\right)}{(k+1)!} \left(\frac{1}{40}\right)^{k+1}}{\frac{\frac{1}{2} \cdot \left(1 - \frac{1}{2}\right) \cdots \left(k - \frac{1}{2} - 1\right)}{k!} \left(\frac{1}{40}\right)^k} \\ &= \frac{k - \frac{1}{2}}{40(k+1)} \leq 1, \end{aligned}$$

that is, $b_2 \geq b_3 \geq \cdots \geq 0$, and $\lim_{k \rightarrow \infty} b_k = 0$ by the Squeeze Theorem because

$$0 \leq b_k \leq \frac{\frac{1}{2} \cdot 1 \cdot 2 \cdots (k-1)}{k!} \left(\frac{1}{40}\right)^k = \frac{1}{2k \cdot 40^k}$$

for $k \geq 2$ and $\lim_{k \rightarrow \infty} \frac{1}{2k \cdot 40^k} = 0$.

By the alternating series estimation, from

$$2 \cdot \left| \binom{\frac{1}{2}}{k+1} \left(\frac{1}{40}\right)^{k+1} \right| < 0.001,$$

we have $k \geq 1$, because $2b_2 = \frac{1}{6400} < 0.001$, and so

$$\sqrt{4.1} \approx 2 + 2 \binom{\frac{1}{2}}{1} \frac{1}{40} = 2.025$$

with error less than 0.001. □

DEFINITION 7.8. A real-valued function f defined on an open interval I is said to be **infinitely differentiable** on I if all (higher) derivatives $f^{(n)}(x)$, $n \geq 1$, exist. The set of infinitely differentiable functions on I is denoted by $C^\infty(I)$.

As a consequence, the functions $f(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$ are infinitely differentiable on $(x_0 - R, x_0 + R)$ if $R > 0$.

COROLLARY 7.9 (Uniqueness Theorem). *Suppose that*

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n \quad \text{and} \quad \sum_{n=0}^{\infty} b_n(x - x_0)^n$$

are two power series which converge for $|x - x_0| < R$ with $R > 0$. Then

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n = \sum_{n=0}^{\infty} b_n(x - x_0)^n \quad \text{for } |x - x_0| < R$$

if and only if

$$a_k = b_k$$

for all $k = 0, 1, 2, 3, \dots$.

PROOF. Suppose that $a_k = b_k$ for all $k = 0, 1, 2, 3, \dots$.

$$\text{Then } \sum_{n=0}^{\infty} a_n(x - x_0)^n = \sum_{n=0}^{\infty} b_n(x - x_0)^n.$$

Conversely, suppose that

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n = \sum_{n=0}^{\infty} b_n(x - x_0)^n = f(x)$$

for $|x - x_0| < R$. Then $a_n = \frac{f^{(n)}(x_0)}{n!}$ and $b_n = \frac{f^{(n)}(x_0)}{n!}$.

Thus $a_n = b_n$ for all $n = 0, 1, 2, 3, \dots$. □

8. Taylor Series

8.1. History Remarks. The study of sequences and series of functions has its origins in the study of power series representation of functions. The power series of $\ln(1 + x)$ was known to Nicolaus Mercator (1620-1687) by 1668, and the power series of many other functions such as $\arctan x$, $\arcsin x$, and etc, were discovered around 1670 by James Gregory (1625-1683). All these series were obtained without any reference to calculus. The first discoveries of Issac Newton (1642-1727), dating back to the early months of 1665, resulted from his ability to express functions in terms of power series. His treatise on calculus, published in 1737, was appropriately entitled *A treatise of the methods of fluxions and infinite series*. Among his many accomplishments, Newton derived the power series expansion of $(1 + x)^{m/n}$ using algebraic techniques. This series and the geometric series were crucial in many of his computations. Newton also displayed the power of his calculus by deriving the power series expansion of $\ln(1 + x)$ using term-by-term integration of the expansion of $1/(1 + x)$. Colin Maclaurin (1698-1746) and Brooks Taylor (1685-1731) were among the first mathematicians to use Newton's calculus in determining the coefficients in the power series expansion of a function. Both realized that if a function $f(x)$ had a power series expansion $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, then the coefficients a_n had to be given by $\frac{f^{(n)}(x_0)}{n!}$.

Why does one care? You probably like polynomials. Think of power series as “generalized” polynomials. Since (almost) all functions you encounter have a Taylor series, all functions can be thought of as “generalized” polynomials!

For instance, you will see that power series are easy to differentiate and integrate. No more techniques of integration, if one is satisfied with writing an integral as a power series!

In finding integrals and solving differential equations, one often faces the problem that the solutions can't be “found”, just because they do not have a name, i.e., they cannot be written down by combining the familiar function names and the familiar mathematical notation. The error function and the functions describing the motion of a “simple” pendulum are important examples. Power series open the door to explore even functions like these!

8.2. Taylor Polynomials and Taylor Series.

DEFINITION 8.1. Let $f(x)$ be a function defined on an open interval I , and let $x_0 \in I$ and $n \geq 1$. Suppose that $f^{(n)}(x)$ exists for all $x \in I$. The polynomial

$$T_n(f, x_0)(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$

is called the **Taylor polynomial of order n** of f at the point x_0 . If f is infinitely differentiable on I , the power series

$$T(f, x_0)(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

is called the **Taylor series** of f and x_0 .

For the special case $x_0 = 0$, the Taylor series of a function f is often referred to as **Maclaurin series**.

The first few Taylor polynomials are as follows:

$$T_0(f, x_0)(x) = f(x_0),$$

$$T_1(f, x_0)(x) = f(x_0) + f'(x_0)(x - x_0),$$

$$T_2(f, x_0)(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2,$$

$$T_3(f, x_0)(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \frac{f'''(x_0)}{3!}(x - x_0)^3.$$

The Taylor polynomial $T_1(f, x_0)$ is the **linear approximation** of f at x_0 , that is **the tangent line** passing through $(x_0, f(x_0))$ with slope $f'(x_0)$.

In general, the Taylor polynomial T_n of f is a polynomial of degree less than or equal to n that satisfies the conditions

$$T_n^{(k)}(f, x_0)(x_0) = f^{(k)}(x_0)$$

for $0 \leq k \leq n$. Since $f^{(n)}(x_0)$ might be zero, T_n could very well be a polynomial of degree strictly less than n .

EXAMPLE 8.2. Find the Maclaurin series of e^x .

SOLUTION. Let $f(x) = e^x$. Then $f^{(n)}(x) = e^x$. Thus $f^{(n)}(0) = 1$ and so the Maclaurin series of e^x is

$$1 + x + \frac{x^2}{2!} + \cdots .$$

□

EXAMPLE 8.3. Find the Taylor series of $f(x) = \sin x$ at $x_0 = \pi$.

SOLUTION.

$$f(x) = \sin x \quad f'(x) = \cos x$$

$$f''(x) = -\sin x \quad f'''(x) = -\cos x \quad \cdots$$

$$f(\pi) = 0 \quad f'(\pi) = -1 \quad f''(\pi) = 0 \quad f'''(\pi) = 1,$$

$$T_3(f, \pi)(x) = 0 - (x - \pi) + 0 + \frac{1}{3!}(x - \pi)^3.$$

$$T(f, \pi)(x) = -(x - \pi) + \frac{1}{3!}(x - \pi)^3 - \frac{1}{5!}(x - \pi)^5 + \cdots$$

$$= \sum_{k=0}^{\infty} (-1)^{k+1} \frac{1}{(2k+1)!} (x - \pi)^{2k+1}.$$

□

EXAMPLE 8.4. Find the Taylor series of $f(x) = \frac{1}{x}$ at $x_0 = 3$.

PROOF.

$$\begin{aligned}\frac{1}{x} &= \frac{1}{3 - (3 - x)} = \frac{1}{3} \cdot \frac{1}{1 - \frac{3-x}{3}} \\ &= \frac{1}{3} \sum_{n=0}^{\infty} \left(\frac{3-x}{3} \right)^n = \frac{1}{3} \sum_{n=0}^{\infty} \frac{(-1)^n}{3^n} (x-3)^n \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n}{3^{n+1}} (x-3)^n.\end{aligned}$$

Thus the Taylor series of $\frac{1}{x}$ at $x_0 = 3$ is

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{3^{n+1}} (x-3)^n.$$

□

8.3. Taylor Theorem. In view of Theorem 7.4, we may ask the following question:

Question: Given an infinitely differentiable function $f(x)$, does the equality

$$(8) \quad f(x) = T(f, x_0)(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$

hold for $|x - x_0| < R$?

(Here R is the radius of the convergence of the Taylor series.)

It turns out that in general, the answer is NO. (See Example 8.5 for an example of a function such that equation 8 does not hold.)

However, as we have seen, the above equality does hold for some elementary functions such as e^x , $\sin x$, $\cos x$, $\ln(1 + x)$, $(1 + x)^a$, $\arctan x$, and etc. We are going to give certain hypothesis such that the above equality holds for some functions.

EXAMPLE 8.5 (Counter-example to the Question). Consider the function

$$f(x) = \begin{cases} e^{-\frac{1}{x^2}} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Then we will show that $f(x) \neq$ its Taylor series at $x_0 = 0$.

PROOF. First we compute $f'(0)$. By substituting $y = \frac{1}{x^2}$

$$\begin{aligned} f'(0) &= \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{e^{-\frac{1}{x^2}} - 0}{x} \\ &= \lim_{x \rightarrow 0} \frac{1}{x e^{\frac{1}{x^2}}} = \lim_{x \rightarrow 0} \frac{1}{x^2 e^{\frac{1}{x^2}}} \cdot x \\ &= \lim_{y \rightarrow \infty} \frac{y}{e^y} \cdot \lim_{x \rightarrow 0} x = 0 \cdot 0 \quad (\text{by L'Hospital's rule}) = 0. \end{aligned}$$

For $x \neq 0$,

$$f'(x) = \frac{d}{dx} \left(e^{-\frac{1}{x^2}} \right) = 2x^{-3} e^{-\frac{1}{x^2}}.$$

Thus,

$$f'(x) = \begin{cases} 2x^{-3} e^{-\frac{1}{x^2}} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Next we compute $f''(x)$. By substituting $y = \frac{1}{x^2}$,

$$\begin{aligned} f''(0) &= \lim_{x \rightarrow 0} \frac{f'(x) - f'(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{2x^{-3} e^{-\frac{1}{x^2}} - 0}{x} \\ &= 2 \lim_{x \rightarrow 0} \frac{1}{x^4 e^{\frac{1}{x^2}}} \end{aligned}$$

$$= 2 \lim_{y \rightarrow \infty} \frac{y^2}{e^y} = 0 \quad (\text{by L'Hospital's rule}).$$

Again, for $x \neq 0$,

$$f''(x) = \frac{d}{dx} \left(2x^{-3} e^{-\frac{1}{x^2}} \right) = (-6x^{-4} + 4x^{-6}) e^{-\frac{1}{x^2}}.$$

Similar calculations will lead to

$$f(0) = f'(0) = f''(0) = f^{(3)}(0) = f^{(4)}(0) = \dots = 0.$$

Thus we have

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k = \sum_{k=0}^{\infty} \frac{0}{k!} x^k = 0 + 0x + 0x^2 + \dots = 0.$$

Clearly, at any $x \neq 0$, $f(x) = e^{-\frac{1}{x^2}} \neq 0$. Therefore,

$$f(x) \neq \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k. \quad \square$$

The **remainder** or **error function** between $f(x)$ and $T_n(f, x_0)$ is defined by

$$R_n(f, x_0)(x) = f(x) - T_n(f, x_0)(x).$$

Clearly

$$\begin{aligned} f(x) &= \lim_{n \rightarrow \infty} T_n(f, x_0)(x) = \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k \\ &= \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k \end{aligned}$$

if and only if

$$\lim_{n \rightarrow \infty} R_n(f, x_0)(x) = 0.$$

To emphasize this fact, we state it as a theorem

THEOREM 8.6. *Suppose that f is an infinitely differentiable function on an open interval I and $x_0 \in I$. Then, for $x \in I$,*

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

if and only if

$$\lim_{n \rightarrow \infty} R_n(f, x_0)(x) = 0.$$

□

The remainder $R_n(f, x_0)$ has been studied much and there are various forms of $R_n(f, x_0)$. We only provide one result called **Lagrange Form of the Remainder**, attributed by Joseph Lagrange (1736-1813). But this result sometimes also referred to as Taylor's theorem.

THEOREM 8.7 (Taylor Theorem). *Let f be a function on an open interval I , $x_0 \in I$ and $n \in \mathbb{N}$. If $f^{(n+1)}(t)$ exists for every $t \in I$, then for any $x \in I$, there exists a ξ between x_0 and x such that*

$$(9) \quad R_n(f, x_0)(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x - x_0)^{n+1}.$$

Thus

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!} (x - x_0)^2 + \cdots \\ + \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n + \frac{f^{(n+1)}(\xi)}{(n+1)!} (x - x_0)^{n+1},$$

for some ξ between x and x_0 .

PROOF. Omitted

□

EXAMPLE 8.8. *Show that, for $|x| < \infty$,*

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots.$$

PROOF. First the right hand side is the Maclaurin series of $f(x) = \sin x$ because

$$f(x) = \sin x \quad f'(x) = \cos x$$

$$f''(x) = -\sin x \quad f'''(x) = -\cos x \quad \cdots$$

$$f(0) = 0 \quad f'(0) = 1 \quad f''(0) = 0 \quad f'''(0) = -1, \quad \cdots$$

Next we show that the remainder tends to 0. Since $f(x) = \sin x$, the higher derivatives $f^{(n+1)}(x)$ is either $\pm \sin x$ or $\pm \cos x$. Thus $|f^{(n+1)}(x)| \leq 1$ for all x and all n .

By the Taylor Theorem,

$$|R_n(f, 0)(x)| = \left| \frac{f^{(n+1)}(\xi)}{(n+1)!} x^{n+1} \right| \leq \frac{|x|^{n+1}}{(n+1)!}.$$

Since $\lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{(n+1)!} = 0$, we have $\lim_{n \rightarrow \infty} |R_n(f, 0)(x)| = 0$, by the Squeeze Theorem, or

$$\lim_{n \rightarrow \infty} R_n(f, 0)(x) = 0$$

for all x . Hence, for all $x \in \mathbb{R}$,

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots.$$

□

8.4. Some Standard Power Series. Below is a list of Maclaurin series of some elementary functions.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots \quad (|x| < \infty)$$

$$\begin{aligned} \sin x &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \\ &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots \quad (|x| < \infty). \end{aligned}$$

$$\begin{aligned} \cos x &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \quad (|x| < \infty). \end{aligned}$$

$$\begin{aligned} \ln(1+x) &= \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{n} \\ &= x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots \quad (-1 < x \leq 1). \end{aligned}$$

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \cdots \quad (|x| < 1).$$

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n = 1 - x + x^2 - x^3 + \cdots \quad (|x| < 1).$$

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$$

$$= x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \quad (-1 \leq x \leq 1)$$

$$(1+x)^a = \sum_{n=0}^{\infty} \binom{a}{n} x^n$$

$$= 1 + \binom{a}{1}x + \binom{a}{2}x^2 + \dots \quad (|x| < 1),$$

where

$$\binom{a}{k} = \frac{a \cdot (a-1) \cdot (a-2) \cdots (a-k+1)}{k!}$$

for any real number a and integers $k \geq 1$, and $\binom{a}{0} = 1$

Remark. From these power series, we can obtain Maclaurin series of various more complicated functions by using operations such as substitution, addition, subtraction, multiplication, division, integrals, derivatives and etc. For the Maclaurin series of $\sin x^2$ can be obtained by replacing x by x^2 in the Maclaurin series of $\sin x$. By using multiplication, we can obtain the Maclaurin series of $e^x \cdot \sin x$. By using long division, we can obtain the Maclaurin series of $\tan x = \frac{\sin x}{\cos x}$. By taking integral, we can obtain the Maclaurin series of non-elementary functions, for instance $f(x) = \int_0^x \sin t^2 dt$.

Example. We can also use power series to define elementary functions. As an example, we start with the exponential function. For any complex number z , let

$$e^z = 1 + z + \frac{z^2}{2!} + \cdots = \sum_{n=0}^{\infty} \frac{z^n}{n!},$$

which is absolutely convergent for any z . You can check the formula

$$\begin{aligned} e^{z_1+z_2} &= \sum_{n=0}^{\infty} \frac{(z_1 + z_2)^n}{n!} = e^{z_1} \cdot e^{z_2} \\ &= \left(\sum_{n=0}^{\infty} \frac{z_1^n}{n!} \right) \cdot \left(\sum_{n=0}^{\infty} \frac{z_2^n}{n!} \right). \end{aligned}$$

For any real number x ,

$$\begin{aligned} e^{ix} &= \sum_{n=0}^{\infty} \frac{(ix)^n}{n!} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!} + i \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!} \\ &= \cos x + i \sin x, \end{aligned}$$

which is the Euler formula.

Computation of π

As an application, we are going to compute number π .

Step 1. Find the Maclaurin series of $\arcsin x$ for $|x| < 1$.

For $|x| < 1$,

$$\begin{aligned}
 \arcsin x &= \int_0^x \frac{1}{\sqrt{1-t^2}} dt = \int_0^x (1 + (-t^2))^{-\frac{1}{2}} dt \\
 &= \int_0^x \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-t^2)^k dt \\
 &= \sum_{k=0}^{\infty} \int_0^x \binom{-\frac{1}{2}}{k} (-1)^k t^{2k} dt \\
 &= \sum_{k=0}^{\infty} (-1)^k \binom{-\frac{1}{2}}{k} \frac{x^{2k+1}}{2k+1} \\
 &= x + \sum_{k=1}^{\infty} (-1)^k \binom{-\frac{1}{2}}{k} \frac{x^{2k+1}}{2k+1}
 \end{aligned}$$

Note that

$$\begin{aligned} \binom{-\frac{1}{2}}{k} &= \frac{\left(-\frac{1}{2}\right) \cdot \left(-\frac{1}{2} - 1\right) \cdots \left(-\frac{1}{2} - k + 1\right)}{k!} \\ &= \frac{\left(-\frac{1}{2}\right) \cdot \left(-\frac{3}{2}\right) \cdots \left(-\frac{2k-1}{2}\right)}{k!} \\ &= (-1)^k \frac{1 \cdot 3 \cdots (2k-1)}{k! \cdot 2^k} \end{aligned}$$

for $k \geq 1$.

Thus

(10)

$$\begin{aligned} \arcsin x &= x + \sum_{k=1}^{\infty} (-1)^k (-1)^k \frac{1 \cdot 3 \cdots (2k-1)}{2^k \cdot k! \cdot (2k+1)} x^{2k+1} \\ &= x + \sum_{k=1}^{\infty} \frac{1 \cdot 3 \cdots (2k-1)}{2^k \cdot k! \cdot (2k+1)} x^{2k+1} \end{aligned}$$

for $|x| < 1$.

Step 2. Find a series expansion of $\frac{\pi}{6}$ using

$$\frac{\pi}{6} = \arcsin \frac{1}{2}.$$

From Equation (10), we obtain the following formula.

$$(11) \quad \frac{\pi}{6} = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (2k-1)}{k! \cdot (2k+1) \cdot 2^{3k+1}}.$$

Step 3. Estimate the remainder.

The remainder of the formula (11) can be estimated as follows.

$$\begin{aligned}
 R_n = |S - S_n| &= \sum_{k=n+1}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (2k-1)}{2^k \cdot k! \cdot (2k+1) \cdot 2^{2k+1}} \\
 &< \sum_{k=n+1}^{\infty} \frac{1}{(2k+1)2^{2k+1}} \\
 &< \sum_{k=n+1}^{\infty} \frac{1}{(2n+3)2^{2k+1}} \\
 &= \frac{1}{(2n+3)2^{2n+3}} \left(1 + \frac{1}{4} + \frac{1}{4^2} + \dots \right) \\
 &= \frac{1}{(2n+3)2^{2n+3} \left(1 - \frac{1}{4}\right)} = \frac{1}{3(2n+3)2^{2n+1}}
 \end{aligned}$$

For instance, let $n = 10$, we have

$$\pi \approx 6 \left(\frac{1}{2} + \sum_{k=1}^{10} \frac{(1 \cdot 3 \cdot \dots \cdot (2k-1))}{k! \cdot (2k+1) \cdot 2^{3k+1}} \right)$$

with error less than

$$6 \cdot \frac{1}{3 \cdot 23 \cdot 2^{21}} = \frac{1}{23 \cdot 2^{20}} = \frac{1}{24117248}.$$

If we choose $n = 20$, we have

$$\pi \approx 6 \left(\frac{1}{2} + \sum_{k=1}^{20} \frac{(2k-1)!!}{k! \cdot (2k+1) \cdot 2^{3k+1}} \right)$$

with error less than

$$6 \cdot \frac{1}{3 \cdot 43 \cdot 2^{41}} = \frac{1}{43 \cdot 2^{40}} = \frac{1}{47278999994368} < 10^{-13}.$$

If $n = 40$, the error is less than

$$\frac{1}{100340843028014221500612608} < 10^{-26}.$$

Remark. There are several other methods for computing π . For instance, we can also use

$$\frac{\pi}{4} = \arctan 1 = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots,$$

but one needs a huge number of terms to get enough accuracy. (So this method is no good for computational purpose!) Another method is to use the formula of John Machin (1680-1751):

$$4 \arctan \frac{1}{5} - \arctan \frac{1}{239} = \frac{\pi}{4},$$

see our text book [6, Problem 7, p.813] for details. Machin used his method in 1706 to find π correct to 100 decimal places. In 1995 Jonathan and Peter Borwein of Simon Fraser University and Yasumasa Kanada of the University of Tokyo calculated the value of π to 4, 294, 967, 286 decimal places!

Another story on computing π is a Chinese mathematician and astronomer Tsu Ch'ung Chi (430-501). He gave the rational approximation $\frac{355}{113}$ to π which is correct to 6 decimal places. He also proved that

$$3.1415926 < \pi < 3.1415927$$

a remarkable result (**Note.** He was a person lived 1500 years ago!), on which it would be nice to have more details but Tsu Ch'ung Chi's book, written with his son, is lost. (His method is to cut off the circle by equal pieces to get his approximation to π .) Tsu's astronomical achievements include the making of a new calendar in 463 which never came into use. (According to the article of J J O'Connor and E F Robertson in <http://www-groups.dcs.st-and.ac.uk/history/Mathematicians/Tsu.html>.)

Remark. Those, who are interested in more applications of Taylor series, can try to finish the applied project, *Radiation from the Stars*, in our text book [6, pp.808-809].

Bibliography

- [1] R. G. Bartle and D. R. Sherbert, *Introduction to real analysis*, 3rd edition, John Wiley, 2000. (Compulsory reading)
- [2] Watson Fulks, *Advanced Calculus*, 3rd Edition, John Wiley & Sons, Inc. press, 1961.
- [3] Wilfred Kaplan, *Advanced Calculus*, 3rd Edition, Addison-Wesley Publishing Company press, 1984.
- [4] William R. Parzynski and Philip W. Zipse, *Introduction to mathematical analysis*, International Edition, McGraw-Hill Book Company press, 1987.
- [5] W. Rudin, *Principles of Mathematical Analysis*, 3rd edition, McGraw-Hill, 1976.
- [6] James Stewart, *Calculus*, 4th Edition, Brooks/Cole Publishing Company press, 1999.
- [7] Manfred Stoll, *Real Analysis*, 2nd Edition, Addison Wesley Longman, Inc. press, 2001.
- [8] G. B. Thomas, Jr. and Ross L. Finney, *Calculus and analytic geometry*, 9th Edition, International Student Edition, Addison-Wesley Longman Inc. press, 1996.